

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶: C07K 14/705, C12N 15/12, A61K 38/17	A1	(11) International Publication Number: WO 96/20957 (43) International Publication Date: 11 July 1996 (11.07.96)
(21) International Application Number: PCT/US95/17095 (22) International Filing Date: 28 December 1995 (28.12.95) (30) Priority Data: 08/366,953 30 December 1994 (30.12.94) US 08/484,397 7 June 1995 (07.06.95) US (71) Applicants: AMGEN INC. [US/US]; Amgen Center, 1840 Dehavilland Drive, Thousand Oaks, CA 91320-1789 (US). THE ROCKEFELLER UNIVERSITY [US/US]; 1230 York Avenue, New York, NY 10021-6399 (US). (72) Inventors: JUAN, Shao, Chieh; 11373 Broadview Drive, Moorpark, CA 93021 (US). LICHENSTEIN, Henri, S.; 9586 Lucerne Street, Ventura, CA 73004 (US). WRIGHT, Samuel, D.; 2 Briar Close, Larchmont, NY 10538 (US). (74) Agents: ODRE, Steven, M. et al.; Amgen Inc., Amgen Center, 1840 Dehavilland Drive, Thousand Oaks, CA 91320-1789 (US).		(81) Designated States: AM, AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LT, LU, LV, MD, MG, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TT, UA, UG, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: ANTI-INFLAMMATORY CD14 POLYPEPTIDES (57) Abstract The invention relates to anti-inflammatory polypeptides comprising soluble CD14 related polypeptides having amino acids at position 7-10 that are different from the native sequence or having amino acids 1-14 deleted.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

- 1 -

ANTI-INFLAMMATORY CD14 POLYPEPTIDES

FIELD OF THE INVENTION

Generally, the invention relates to the field
5 of polypeptides that have anti-inflammatory properties.
Amino acids 7 to 10 of CD14 have been found to contain
an important domain enabling inflammatory responses in
cells, including IL-6 production. The polypeptides of
this invention were based on replacing amino acids
10 7 - 10 in the soluble form of CD14 with different amino
acids. Other polypeptides of this invention are missing
the first 14 amino acids of soluble CD14. The
polypeptides of the invention may be used to treat
inflammatory conditions, such as sepsis.

15

BACKGROUND OF THE INVENTION

Sepsis is a life-threatening medical condition
that can be brought on by infection or trauma. The
symptoms of sepsis can include chills, profuse sweating,
20 fever, weakness, or hypotension, followed by leukopenia,
intravascular coagulation, shock, adult respiratory
distress syndrome, multiple organ failure, and often,
death. R. Ulevitch, et al., J. Trauma 30: S189-92
(1990).

25

The lipopolysaccharides ("LPS"; also,
"endotoxins") that are typically present on the outer
membrane of all gram-negative bacteria are among the
most studied and best understood sepsis-inducing
substances. While the precise chemical structures of
30 LPS molecules obtained from different bacteria may vary
in a species-specific fashion, a region called the lipid
A region is common to all LPS molecules. E. Rietschel
et al., in Handbook of Endotoxins, 1: 187-214, eds. R.
Proctor and E. Rietschel, Elsevier, Amsterdam (1984).

- 2 -

This lipid A region is responsible for many, if not all, of the LPS-dependent pathophysiologic changes that characterize sepsis.

LPS is believed to be a primary cause of death
5 in humans afflicted with gram-negative sepsis. van
Deventer et al., Lancet, 1: 605 (1988); Ziegler et al.,
J. Infect. Dis., 136: 19-28 (1987). Treatment of
patients suffering from sepsis and gram-negative
bacteraemia with a monoclonal antibody against LPS
10 decreased their mortality rate. Ziegler et al.,
N. Eng. J. Med., 324: 429 (1991).

Sepsis is also caused by gram-positive
bacteria. Bone, R.C. Arch. Intern. Med., 154: 26-34
(1994). The activation of host cells can originate from
15 gram-positive cell walls or purified cell components
such as peptidoglycan and lipoteichoic acid. Such
substances induce a similar pattern of inflammatory
responses to those induced by LPS. Chin and Kostura,
J. Immunol. 151: 5574-5585 (1993); Mattson et al., FEMS
20 Immun. Med. Microbiol. 7: 281-288 (1993); and Rotta,
J. Z. Immunol. Forsch. Bd. 149: 230-244 (1975).

LPS and gram-positive cell wall substances
cause polymorphonuclear leukocytes, endothelial cells,
and cells of the monocyte/macrophage lineage to rapidly
25 produce and release a variety of cell products,
including cytokines, which are capable of initiating,
modulating or mediating humoral and cellular immune
responses and processes.

One particular cytokine, alpha-cachectin or
30 tumor necrosis factor (TNF- α), is apparently a primary
mediator of septic shock. Beutler et al., N. Eng. J.
Med., 316: 379 (1987). Intravenous injection of LPS
into experimental animals and man produces a rapid,
transient release of TNF- α . Beutler et al., J.
35 Immunol., 135: 3972 (1985); Mathison et al., J. Clin.
Invest. 81: 1925 (1988). Pretreatment of animals with

- 3 -

anti-TNF- α antibodies can modulate septic shock.

Beutler et al., Science, **229**: 869, (1985); Mathison et al., J. Clin. Invest. **81**: 1925 (1988).

5 Molecular receptors that can combine with sepsis-inducing substances, and that once combined, initiate certain chemical reactions, play a critical role in the etiology of the symptoms of sepsis. CD14 is a 55-kD glycoprotein expressed strongly on the surface of monocytes and macrophages, and weakly on the surface
10 of granulocytes, such as neutrophils. S. M. Goyert et al., J. Immunol. **137**: 3909 (1986). A. Haziot et al., J. Immunol. **141**: 547-552 (1988); S. M. Goyert et al., Science **239**: 497 (1988).

The cDNAs and the genes for human and murine
15 CD14 have been cloned and sequenced. E. Ferrero and S.M. Goyert, Nuc. Acids Res. **16**: 4173 (1988); S. M. Goyert et al., Science **239**: 497 (1988); M. Setoguchi et al., Biochem. Biophys. Acta **1008**: 213-22 (1989). CD14 is linked by a cleavable glycosyl phosphatidyl inositol
20 tail [A. Haziot et al., J. Immunol. **141**: 547-552 (1988)] to the exoplasmic surface of mature monocytes, macrophages, granulocytes and dendritic reticulum cells, or renal nonglomerular endothelium, and of hepatocytes in rejected livers.

25 CD14 mediates responses by binding to LPS. Complexes of LPS and sCD14 exhibit a 1:1 stoichiometry (Hailman, E., et al., J. Exp. Med. **179**: 269-277 (1994)), and these complexes initiate TNF- α production in monocytes (Dentener, M.A., et al., J. Immunol. **7**: 2885-
30 2891 (1993)), IL-6 production in astrocytes (Frey, E., et al., J. Exp. Med. **176**: 1665-1671 (1992)), production of adhesion molecules in endothelial cells (Frey, E., et al., J. Exp. Med. **176**: 1665-1671 (1992)) and activation of leukocyte integrins in PMN (Hailman, E., et al., J. Exp. Med. **179**: 269-277 (1994)). Spontaneous binding of
35 LPS to CD14 is slow, but this binding may be

- 4 -

dramatically accelerated by LBP. LBP acts in a catalytic fashion, with one molecule of LBP transferring hundreds of LPS molecules to hundreds of CD14 molecules.

Other experiments have shown that cell
5 activation can also be induced by interaction of CD14 with components of gram-positive bacteria such as *B. subtilis*, *S. aureus*, and *S. mitis* (Pugin et al., Immunity 1: 509-516 (1994)). Furthermore, interaction of
10 CD14 with lipoarabinomannan from the cell wall of *Mycobacterium tuberculosis* also induces cellular activation in a CD14-dependent fashion (Zhang et al., J. Clin. Invest. 91: 2076-2083 (1993); Pugin et al., Immunity 1: 509-516 (1994)). These studies suggest that
15 CD14 is a receptor which recognizes a wide variety of bacterial structures. Interaction of CD14 with these structures initiates host inflammatory responses.

Several neutralizing monoclonal antibodies (mAbs) to CD14 have been shown to antagonize cellular responses to LPS in vitro (Wright, S. D., et al. Science
20 249: 1431-1433 (1990); Hailman, E., et al. J. Exp. Med. 179, 269-277 (1994); Frey, E. A., et al. J. Exp. Med. 176, 1665-1671 (1992); Arditi, M., et al. Infect. Immun. 61, 3149-3156 (1993); Wright, S. D., et al. J. Exp. Med. 173, 1281-1286 (1991); Dentener, M. A., et al.
25 J. Immunol 150, 2885-2891 (1993); Grunwald, U., et al. J. Immunol Methods 155, 225-232 (1992)) and in vivo (Leturcq, D. J., et al. Satellite Meeting of the 3rd Conference of the International Endotoxin Society 22
30 (Abstract) (1994)); and Wright, et al., Science 90:1431-1433 (1990). Additional in vivo data have demonstrated that animals injected with CD14 neutralizing monoclonal antibodies become hyporesponsive to LPS and mice lacking CD14 fail to respond to LPS. These experiments suggest
35 that the release of inflammatory cytokines can be blocked by preventing the interaction of LPS with membrane CD14.

- 5 -

CD14 has also been shown to exist as a soluble protein found in normal sera or urine of nephrotic patients. Recent evidence has shown that sCD14 enables LPS-dependent responses in cells which lack membrane

5 CD14, i.e., endothelial cells and epithelial cells. In these cells types, sCD14 in conjunction with LPS promotes inflammatory cytokine release and upregulation of adhesion molecules.

Interestingly, high concentrations of sCD14

10 have been shown to block inflammatory cytokine release from monocytes in a whole blood assay. Presumably, the beneficial effect of sCD14 in this assay arises from its ability to divert LPS away from mCD14 on macrophages and PMNs. Thus sCD14, like CD14 neutralizing monoclonal

15 antibodies, could be useful in preventing LPS interactions in mCD14. However, the utility of sCD14 to treat LPS-mediated inflammatory disorders is limited by its other property of eliciting inflammatory cytokines in endothelial cells. Thus, a sCD14 molecule which

20 retained its ability to bind LPS, yet did not activate endothelial cells should have superior properties in treating inflammation.

Monoclonal antibodies may be a useful tool to help identify domains in sCD14 required for cell

25 activation. We have demonstrated that mAbs MEM-18 and 3C10 recognize a sCD14 mutant truncated at amino acid 152, indicating that epitopes for these two mAbs are within the first 152 amino acids (Juan, T. S. -C., et al. J. Biol. Chem. **270**, 1382-1387 (1995)). We further

30 localized the epitope of MEM-18 between amino acids 57 and 64 and found that this region is also essential for LPS binding (Juan, T. S. -C., et al. J. Biol. Chem. **270**, 5219-5224 (1995)). Deletion of this region not only disrupted binding of MEM-18, but also binding of LPS.

35 The epitope for mAb 3C10 defines another functional domain of CD14. This mAb appears to

- 6 -

recognize a different region from that of MEM-18 (Juan, T. S. -C., et al. J. Biol. Chem. 270, 1382-1387 (1995)). Binding of monoclonal antibody 3C10 to sCD14 does not affect LPS binding to sCD14 (Juan, T. S. -C., et al. J. Biol. Chem. 270, 5219-5224 (1995)), suggesting that this epitope may be involved in a cellular function other than LPS binding.

For the preceding reasons, it is an object of this invention to develop methods and therapies for the effective treatment, including prevention, for symptoms of inflammatory conditions, including sepsis. It is also an object of this invention to develop methods and therapies for the effective protection of individuals who are at risk of becoming afflicted by the symptoms of inflammation, including sepsis.

It is another object of this invention to develop methods and therapies for the effective treatment, including prevention, of symptoms of diseases that are mediated by LPS, gram-negative bacteraemia, gram-positive cell components, gram-positive bacteraemia, mycobacterial lipoarabinomannan, mycobacterial infections and/or CD14. Such diseases include ARDS, septic shock, acute pancreatitis, acute and chronic liver failure, intestinal or liver transplantation, inflammatory bowel disease, graft vs. host disease in bone marrow transplantation and tuberculosis.

SUMMARY OF THE INVENTION

30

The present inventors have discovered a group of polypeptides that are capable of binding to lipopolysaccharide (LPS), resulting in inhibition of the binding of LPS or gram-positive cell components to membrane CD14, thus reducing or eliminating CD14-mediated inflammatory responses. As used herein,

- 7 -

inhibition of binding of LPS also means inhibition of binding to gram-positive cell components. This group of polypeptides was designed by the inventors based on their important discovery of an LPS-binding domain in
5 CD14.

The polypeptides of this invention are based on substituting the naturally-occurring amino acids at positions 7-10 of CD14, preferably in a soluble form, with neutral amino acids, preferably those having either
10 a hydrogen or C1-C6 alkyl side chain. The polypeptides of this invention also include those that begin at amino acid 15 of FIG. 1. With reference to FIG.1, "soluble" CD14 (sCD14) is a molecule selected from sequences starting at an amino acid of from 1 through 6 and ending
15 at an amino acid of from 152 through 348. Native human CD14 has as X₁-X₄, Glu-Leu-Asp-Asp in FIG.1. Preferably, the polypeptide is substituted with alanine at each of positions 7 through 10.

The polypeptides of this invention are capable
20 of binding to LPS, thereby preventing further binding of microbial cell components to membrane CD14. If microbial cell interaction with membrane CD14 is prevented, the cascade of events leading to inflammation, and especially sepsis, are reduced or
25 prevented. Therefore, the polypeptides of this invention have anti-inflammatory properties.

More specifically, the evidence provided herein indicates, inter alia, that a polypeptide having the region from amino acids 7 to 10 in CD14 replaced
30 with alanine residues or having amino acids 1 to 14 deleted from soluble CD14, binds LPS but mediates a substantially reduced cellular inflammatory response, such as production of the cytokine IL-6 in response to LPS, as compared to native CD14.

35 In the examples below, we identify the epitope of 3C10 by making a series of site-directed alanine

- 8 -

substitution mutants in sCD14. We show that the region between amino acids 7 and 14 are required for 3C10 binding. We further characterized this domain by generating a sCD14 mutant with alanine substituted at amino acids 7 to 10 (sCD14_{(7-10)A}). This mutant was capable of binding LPS but was impaired in its ability to mediate cellular responses to LPS.

The peptides and polypeptides of this invention may be prepared by (a) standard synthetic methods, (b) derivation from CD14, (c) recombinant methods, (d) a combination of one or more of (a) - (c), or other methods of preparing polypeptides.

The polypeptides of this invention may be used for therapeutic or prophylactic purposes by incorporating them into appropriate pharmaceutical carrier materials and administering an effective amount to a patient, such as a human (or other mammal) in need thereof.

BRIEF DESCRIPTION OF THE FIGURES

Numerous other aspects and advantages of the present invention will therefore be apparent upon consideration of the following detailed description thereof, reference being made to the drawings wherein:

FIG. 1 shows a schematic map of soluble human CD14 having 348 amino acids; amino acids 7 - 10 (shown as X₁-X₄) are Glu-Leu-Asp-Asp in native mature human CD14. In the polypeptides of this invention, amino acids 7 - 10 may be any of the amino acids described herein.

FIG. 2 shows the sequence and expression of sCD14 alanine-substitution mutants.

FIG. 3 shows the results of a BIAcore analysis of the monoclonal antibody 3C10 binding to alanine substitution mutants of sCD14. Conditioned media (CM)

- 9 -

were collected from COS-7 cells transfected with no DNA (MOCK), sCD14₁₋₃₄₈, or sCD14 mutants four days after electroporation. All CM were analyzed for their ability to bind the antibody 3C10 as described in the Examples section herein. Relative response units (RRU) were recorded from four repeats of one experiment and calculated as means \pm standard deviations.

FIG. 4 shows that mAb 3C10 does not recognize purified sCD14_{(7-10)A}. Immobilization of mAb 3C10 to a sensor chip has been described (Juan, T. S. -C., et al. J. Biol. Chem. 270, 1382-1387 (1995)). 10 μ g/ml sCD14₁₋₃₄₈ or sCD14_{(7-10)A} was used for injection. Injection of solutions at various "Steps" are marked on the sensorgram. "Wash" indicates a washing step using HBS buffer as described in the Example Section. The experiments were performed three times and the results of one experiment are shown.

FIG. 5 shows that sCD14_{(7-10)A} is defective in enabling cellular responses to LPS. A. sCD14_{(7-10)A} has reduced ability to stimulate IL-6 production by U373 cells. U373 cells were treated with various concentrations of sCD14₁₋₃₄₈ or sCD14_{(7-10)A} in the presence or absence of LPS (20 ng/ml) for 24 h. IL-6 levels were determined as described (Juan, T. S. -C., et al. J. Biol. Chem. 270, 1382-1387 (1995)). Data presented are means \pm standard deviations from four readings in an experiment repeated 3 times. B. sCD14₁₋₃₄₈ but not sCD14_{(7-10)A} mediates responses of PMN to LPS and LBP. Freshly isolated PMN were incubated with "smooth" LPS (*E. coli* 0111:B4 30 ng/ml), rLBP (1 μ g/ml), and the indicated concentrations of sCD14₁₋₃₄₈ or sCD14_{(7-10)A} for 10 min at 37°C. Cells were washed and adhesion to fibrinogen-coated wells was measured (Hailman, E., et al. J. Exp. Med. 179, 269-277 (1994),25). Error bars indicate standard deviations of triplicate determinations.

- 10 -

FIG. 6 shows that sCD14_{(7-10)A} does not activate NF- κ B. Whole cell extracts of U373 cells with various treatments (lane 1, control; lane 2, LPS; lane 3, sCD14₁₋₃₄₈; lane 4, sCD14₁₋₃₄₈ and LPS; lane 5, sCD14_{(7-10)A}; lane 6, sCD14_{(7-10)A} and LPS; lane 7, sCD14 Δ ₍₅₇₋₆₄₎; and lane 8, sCD14 Δ ₍₅₇₋₆₄₎ and LPS) were obtained and binding of proteins to the labeled NF- κ B oligonucleotide was performed as described in the Example Section. Complexes of NF- κ B were resolved on a native-4.5% polyacrylamide gel. After electrophoresis, the gel was dried and exposed to X-ray film for 16 h. Complexes of labeled probe and NF- κ B are indicated.

FIG. 7 shows that sCD14_{(7-10)A} forms stable complexes with ³H-LPS. Various concentrations of sCD14₁₋₃₄₈ (lanes 2-4) or sCD14_{(7-10)A} (lanes 5-7) were incubated with 3 μ g/ml ³H-LPS in the absence (A) or presence of 16.7 nM rLBP (B) as described in the Example Section. Lane 1 contains LPS in the absence of additional protein. Mixtures were run on 4-20% native polyacrylamide gels and processed for fluorography. Positions of uncomplexed LPS and complexes between LPS and sCD14₁₋₃₄₈ or sCD14_{(7-10)A} are indicated.

FIG. 8 shows inhibition of LPS-induced cellular responses by sCD14_{(7-10)A}. A. Inhibition of LPS-induced PMN adhesion by sCD14_{(7-10)A}. Rough LPS (*Salmonella minnesota* R60, 10 ng/ml) was incubated with LBP and various concentrations of sCD14₁₋₃₄₈ or sCD14_{(7-10)A} at 37°C for 30 min before addition of PMN. The adhesion of PMN to fibrinogen was measured as described in the Example Section. Error bars indicate standard deviations from three readings. B. Inhibition of TNF- α production in whole blood by sCD14_{(7-10)A}. 250 μ l of whole blood was incubated with various concentrations of bovine serum albumin, sCD14₁₋₃₄₈, or sCD14_{(7-10)A} in the presence of 0.25 ng/ml smooth LPS (*Salmonella minnesota* wildtype) at 37°C for 3h and TNF- α production was

- 11 -

measured as described in the Example Section. Fraction of TNF- α production refers to the ratio of TNF- α produced in the presence of exogenous protein divided by TNF- α produced in the absence of added protein. Error

5 bars are standard deviations from six readings.

FIG. 9 shows that gram positive cell components compete with LPS for binding to sCD14. ^3H -LPS (1 $\mu\text{g}/\text{ml}$) and sCD14 (50 $\mu\text{g}/\text{ml}$) were incubated alone (lane 1) or with LPS (lane 2) or with *Staphylococcus aureus* crude
10 extract (SACE) (lanes 3-6) at 37° C for 17 hours in PBS with 1 mM EDTA. The samples were then run in a native polyacrylamide gel, and the position of radioactive bands was determined by radioautography.

15 DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

The present invention is based on the discovery by the present inventors of a region on CD14 that is involved in inflammatory cellular responses
20 mediated by CD14, such as production of IL-6 in response to gram negative or gram positive bacteria cell components (e.g., LPS from gram negative bacteria). The Examples below explain in detail the evidence supporting these discoveries.

25 FIG. 1 shows a map of human CD14, including the LPS binding and IL-6 inducing region, corresponding to amino acids 7 to 10.

The polypeptides of this invention are based on substituting amino acids at positions 7 - 10,
30 inclusive, in soluble CD14, with amino acids that are different from those in the native molecule or deleting the first 14 amino acids in soluble CD14. Referring to FIG. 1, amino acids 7 - 10 in the polypeptides that contain substituted amino acids, are different from Glu-
35 Leu-Asp-Asp. Preferably, the substituted amino acids are neutral amino acids. The side chains of these

- 12 -

neutral amino acids are preferably selected from the group consisting of hydrogen and alkyl groups having from 1 to 6 carbon atoms, which may be substituted by one or more substituents selected from halogen (e.g.,
5 Cl, Br, I), -OH, -CN, -OR (R= alkyl having 1 to 6 carbon atoms), and related structures. The specific amino acids substituted at amino acids 7 - 10 are relatively unimportant as long as the resulting polypeptide is capable of binding LPS and/or inhibiting release of
10 inflammatory mediators in monocytes and PMNs while at the same time possessing a diminished ability (compared to wild type sCD14) to induce an inflammatory response in endothelial cells and epithelial cells. The polypeptides of this invention that have the initial 14
15 amino acids deleted from soluble CD14 end at an amino acid in sCD14 of from 152 to 348, inclusive. Preferred deletion-type peptides are amino acids 15-152 and 15-348 of FIG. 1.

The remainder of the polypeptide (other than
20 the substituted amino acids) is generally identical to native soluble CD14. "Soluble CD14" (sCD14) means polypeptides that correspond to amino acids of from positions 1 through 6 to positions 152 through 348, of FIG. 1. Note that in all cases herein the numbering of
25 amino acids used herein corresponds to the amino acids sequence of FIG. 1. For example, even if the particular soluble CD14 starts with the second through sixth amino acid as set forth in FIG. 1, the amino acids to be substituted are those corresponding to amino acids
30 7 - 10 of the full sequence of FIG. 1.

For purposes of this disclosure, the molecule depicted in FIG. 1 will be referred to as sCD14 or sCD14₁₋₃₄₈. Other examples of soluble CD14 will be named by providing the beginning and ending amino acids based
35 on the numbering scheme of FIG. 1; e.g., sCD14₁₋₁₅₂; sCD14₂₋₁₅₂; sCD14₁₅₋₁₅₂; sCD14₁₅₋₃₄₈, etc. Polypeptides

- 13 -

having amino acids substituted for those set forth in FIG.1 will be named as follows: sCD14₁₋₃₄₈ having alanine in place of amino acids 7 - 10 in FIG. 1 is named sCD14₁₋₃₄₈(7-10)A, where "A" is the one-letter code for the amino acid alanine. sCD14₁₋₃₄₈ having an alanine at positions 7 and 9-10, and a glycine at position 8 is named sCD14₁₋₃₄₈(7, 9, 10)A(8)G.

Some specific preferred examples of polypeptides of this invention are set forth below:

10	sCD14 ₁₋₃₄₈ (7-10)G	sCD14 ₁₋₁₅₂ (7-10)G
	sCD14 ₁₋₃₄₈ (7-10)A	sCD14 ₁₋₁₅₂ (7-10)A
	sCD14 ₁₋₃₄₈ (7-10)V	sCD14 ₁₋₁₅₂ (7-10)V
	sCD14 ₁₋₃₄₈ (7-10)L	sCD14 ₁₋₁₅₂ (7-10)L
15	sCD14 ₁₋₃₄₈ (7-10)I	sCD14 ₁₋₁₅₂ (7-10)I
	sCD14 ₁₋₃₄₈ (7-10)P	sCD14 ₁₋₁₅₂ (7-10)P
	sCD14 ₁₅₋₃₄₈	sCD14 ₁₅₋₁₅₂

The present invention also encompasses physiologically acceptable salts of the polypeptides disclosed herein. Also, in each polypeptide, one or more D or L amino acids may be included; however, it is preferred that all of the amino acids are of the L stereochemistry.

The polypeptides of this invention are expected to have the ability to reduce inflammatory responses in cells as compared to native sCD14. Reduction in inflammation may conveniently be measured by examining IL-6 production by such cells using, e.g., the method described in Example 4 below. Preferably the amount of IL-6 reduction, as compared to native sCD14, will be at least 5-fold, particularly preferably, at least 10-fold.

Also, in each case, the amino acids may be chemically derivatized as long as LPS binding coupled with reduced ability to induce IL-6 (or related

- 14 -

cytokines) is retained. Thus, "chemical derivatives" of the present polypeptides are included within the scope of the term "polypeptide" as used herein. These chemical derivatives contain additional chemical moieties not part of the X₁ - X₄ amino acid substituted polypeptides.

Covalent modifications of the polypeptides are included within the scope of this invention. Such modifications may be introduced into the molecule by reacting targeted amino acid residues of the polypeptide with an organic derivatizing agent that is capable of reacting with selected side chains or terminal residues.

Histidyl residues are derivatized by reaction with diethylpyrocarbonate at pH 5.5-7.0 because this agent is relatively specific for the histidyl side chain. Para-bromophenacyl bromide also is useful; the reaction is preferably performed in 0.1 M sodium cacodylate at pH 6.0.

Lysiny and amino terminal residues are reacted with succinic or other carboxylic acid anhydrides. Derivatization with these agents has the effect of reversing the charge of the lysiny residues. Other suitable reagents for derivatizing alpha-amino-containing residues include imidoesters such as methyl picolinimidate; pyridoxal phosphate; pyridoxal; chloroborohydride; trinitrobenzenesulfonic acid; O-methylisourea; 2,4 pentanedione; and transaminase-catalyzed reaction with glyoxylate.

Arginy residues are modified by reaction with one or several conventional reagents, among them phenylglyoxal, 2,3-butanedione, 1,2-cyclohexanedione, and ninhydrin. Derivatization of arginine residues requires that the reaction be performed in alkaline conditions because of the high pK_a of the guanidine functional group. Furthermore, these reagents may react with the groups of lysine as well as the arginine epsilon-amino group.

- 15 -

The specific modification of tyrosyl residues per se has been studied extensively, with particular interest in introducing spectral labels into tyrosyl residues by reaction with aromatic diazonium compounds or

5 tetranitromethane. Most commonly, N-acetylimidazole and tetranitromethane are used to form O-acetyl tyrosyl species and 3-nitro derivatives, respectively.

Carboxyl side groups (aspartyl or glutamyl) are selectively modified by reaction with carbodiimides

10 (R'-N-C-N-R') such as 1-cyclohexyl-3-(2-morpholinyl)-(4-ethyl) carbodiimide or 1-ethyl-3-(4-azonia-4,4-dimethylpentyl) carbodiimide. Furthermore, aspartyl and glutamyl residues are converted to asparaginyl and glutaminyl residues by reaction with ammonium ions.

15 Glutaminyl and asparaginyl residues are frequently deamidated to the corresponding glutamyl and aspartyl residues. Alternatively, these residues are deamidated under mildly acidic conditions. Either form of these residues falls within the scope of this invention.

20 Derivatization with bifunctional agents is useful for cross-linking the peptides or their functional derivatives to a water-insoluble support matrix or to other macromolecular carriers. Commonly used cross-linking agents include, e.g., 1,1-bis(diazoacetyl)-2-phenylethane, 25 glutaraldehyde, N-hydroxysuccinimide esters, for example, esters with 4-azidosalicylic acid, homobifunctional imidoesters, including disuccinimidyl esters such as 3,3' - dithiobis (succinimidylpropionate), and bifunctional maleimides such as bis-N-maleimido-1,8-octane. Derivatizing 30 agents such as methyl-3-[(p-azidophenyl)dithio]propioimide yield photoactivatable intermediates that are capable of forming crosslinks in the presence of light. Alternatively, reactive water-insoluble matrices such as cyanogen bromide-activated carbohydrates and the reactive substrates 35 described in U.S. Pat. Nos. 3,969,287; 3,691,016; 4,195,128;

- 16 -

4,247,642; 4,229,537; and 4,330,440 are employed for protein immobilization.

Other modifications include hydroxylation of proline and lysine, phosphorylation of hydroxyl groups of seryl or threonyl residues, oxidation of the sulfur atom in Cys, methylation of the alpha-amino groups of lysine, arginine, and histidine side chains (T. E. Creighton, Proteins: Structure and Molecule Properties, W. H. Freeman & Co., San Francisco, pp. 79-86 (1983)), acetylation of the N-terminal amine, and, in some instances, amidation of the C-terminal carboxyl groups.

The activity of the polypeptide variant can be screened in a suitable screening assay for the desired characteristic. Biological activity is screened in an appropriate bioassay, as described herein. For example, binding of LPS to CD14 (or a polypeptide of this invention) may be measured in a standard competitive binding assay. Activity to reduce cellular inflammatory responses may be measured in terms of reduction of IL-6 production by cells (e.g., U373 cells) as described herein.

Modifications of such polypeptide properties as redox or thermal stability, hydrophobicity, susceptibility to proteolytic degradation or the tendency to aggregate with carriers or into multimers are assayed by methods well known to the ordinarily skilled artisan.

Such derivatized moieties may improve the solubility, absorption, biological half life, and the like. The moieties may alternatively eliminate or attenuate any undesirable side effect of the protein and the like. Moieties capable of mediating such effects are disclosed, for example, in Remington's Pharmaceutical Sciences, 16th ed., Mack Publishing Co., Easton, Pa. (1980).

The polypeptides of the invention may also be covalently or noncovalently associated with a carrier

- 17 -

molecule, such as a polypeptide or non-CD14 protein, a linear polymer (such as polyethylene glycol, polylysine, etc), a branched-chain polymer (see, for example, U.S. Patent 4,289,872 to Denkenwalter et al., issued September 15, 1981; 5,229,490 to Tam, issued July 20, 1993; WO 93/21259 by Frechet et al., published 28 October 1993); a lipid; a cholesterol group (such as a steroid; or a carbohydrate or oligosaccharide.

The polypeptides of this invention are expected to have the ability to bind to LPS. This binding renders LPS unable to bind to membrane CD14 (mCD14) on macrophages and therefore produces an anti-inflammatory response in a mammal. Additionally, the polypeptides of this invention have reduced ability to trigger an inflammatory response in cells lacking mCD14 such as endothelial and epithelial cells. They are also expected to bind to cellular components of gram positive cells that cause inflammation (analogous to LPS; however, the structure(s) in gram positive bacteria that cause inflammatory responses to cells is (are) not yet known).

"Binding" to LPS means that in a standard competition assay, the polypeptide is capable of inhibiting 50% binding of CD14 to LPS between 1 mM and 1 nM, preferably 100 μ M to 10 nM (IC₅₀ values). A standard binding assay may be carried out as is well known in the art.

The polypeptides of this invention may be made in a variety of ways. For example, solid phase synthesis techniques may be used. Suitable techniques are well known in the art, and include those described in Merrifield, in Chem. Polypeptides, pp. 335-61 (Katsoyannis and Panayotis eds. 1973); Merrifield, J. Am. Chem. Soc., 85, 2149 (1963); Davis et al., Biochem. Int'l, 10, 394-414 (1985); Stewart and Young, Solid Phase Peptide Synthesis (1969); U.S. Pat. No. 3,941,763;

- 18 -

Finn et al., in The Proteins, 3rd ed., vol. 2, pp. 105-253 (1976); and Erickson et al. in The Proteins, 3rd ed., vol. 2, pp. 257-527 (1976).

More preferably, the polypeptides are made in
5 transformed host cells using recombinant DNA techniques. To do so, a recombinant DNA molecule coding for the polypeptide is prepared. Methods of preparing such DNA molecules are well known in the art. For instance, sequences coding for the polypeptides could be excised
10 from DNA using suitable restriction enzymes. Alternatively, the DNA molecule could be synthesized using chemical synthesis techniques, such as the phosphoramidite method. Also, a combination of these techniques could be used.

15 The invention also includes a vector capable of expressing the peptides in an appropriate host. The vector comprises the DNA molecule that codes for the peptides operatively linked to appropriate expression control sequences. Methods of effecting this operative
20 linking, either before or after the DNA molecule is inserted into the vector, are well known. Expression control sequences include promoters, activators, enhancers, operators, ribosomal binding sites, start signals, stop signals, cap signals, polyadenylation
25 signals, and other signals involved with the control of transcription or translation.

The resulting vector having the DNA molecule thereon is used to transform an appropriate host. This transformation may be performed using methods well known
30 in the art.

Any of a large number of available and well-known host cells may be used in the practice of this invention. The selection of a particular host is dependent upon a number of factors recognized by the
35 art. These include, for example, compatibility with the chosen expression vector, toxicity to it of the peptides

- 19 -

encoded for by the DNA molecule, rate of transformation, ease of recovery of the peptides, expression characteristics, bio-safety and costs. A balance of these factors must be struck with the understanding that not all hosts may be equally effective for the expression of a particular DNA sequence.

Within these general guidelines, useful microbial hosts include bacteria (such as *E. coli* sp.), yeast (such as *Saccharomyces* sp.) and other fungi, insects, plants, mammalian (including human) cells in culture, or other hosts known in the art.

Next, the transformed host is cultured under conventional fermentation conditions so that the desired peptides are expressed. Such fermentation conditions are well known in the art.

Finally, the polypeptides are purified from the culture. These purification methods are also well known in the art.

The polypeptides of this invention may be used in any of a number of situations where LPS/gram positive cell component binding is required. For example, therapeutically and prophylactically, the polypeptides may be used for inflammatory bowel disease, acute and chronic liver failure, graft vs. host disease (bone marrow transplant), intestinal or liver transplant, ARDS, acute pancreatitis and tuberculosis. Septic shock is a particularly preferred target condition.

The novel polypeptides are useful for the prophylaxis or treatment of septic shock in mammals, including humans, at doses of about 0.1 to 100 mg/kg of body weight, preferably at a level of about 1 to 50 mg/kg of body weight, and the amount may be administered, e.g., in divided doses on daily basis. The polypeptides may be administered prophylactically to patients who may be exposed to or have been exposed to organisms which may cause septic shock or to detoxify

- 20 -

LPS (bacterial endotoxins) by the use of the same dose set forth above *in vivo*; *in vitro* detoxification or prevention of endotoxin contamination may be carried out at a level which is effective to achieve the desired
5 result. The amount may be based on routine experimentation based on the premise that about 1 mole of endotoxin is bound by 1 mole of polypeptide. The particular dose of a particular polypeptide may be varied within or without the range that is specified
10 herein depending on the particular application or severity of a disease and the condition of the host. Those who are skilled in the art may ascertain the proper dose using standard procedures.

The pharmaceutical compositions of the present
15 invention may be administered by any means that achieve their intended purpose. For example, administration may be by parenteral routes, including subcutaneous, intravenous, intradermal, intramuscular, intraperitoneal, intrathecal, transdermal, or buccal routes. Alternatively, or
20 concurrently, administration may be by the oral or rectal route. The pharmaceutical compositions can be administered parenterally by bolus injection or by gradual perfusion over time.

In addition to the polypeptide, these
25 pharmaceutical compositions may contain suitable pharmaceutically acceptable carriers comprising excipients and auxiliaries which facilitate processing of the active compounds into preparations which can be used pharmaceutically. Preferably, the preparations, particularly
30 those which can be administered orally and which can be used for the preferred type of administration, such as tablets, dragees, and capsules, and also preparations which can be administered rectally, such as suppositories, as well as suitable solutions for administration by injection or
35 orally, contain from about 0.1 to about 99 percent,

- 21 -

preferably from about 25-85 percent, of active compound(s), together with the excipient.

Suitable excipients are, in particular, fillers such as sugars, such as lactose, sucrose, mannitol, or sorbitol; cellulose preparations and/or calcium phosphates, such as tricalcium phosphate or calcium hydrogen phosphate; as well as binders such as starch paste made using, for example, maize starch, wheat starch, rice starch, potato starch, gelatin, gum tragacanth, methyl cellulose, hydroxypropylmethylcellulose, sodium carboxymethyl cellulose, and/or polyvinylpyrrolidone. If desired, disintegrating agents may also be added, such as the above-mentioned starches as well as carboxymethyl starch, cross-linked polyvinyl pyrrolidone, agar, or alginic acid or a salt thereof, such as sodium alginate. Auxiliaries which can be used in the compositions according to the present invention include flow-regulating agents and lubricants such as silica, talc, stearic acid or salts thereof, a detergent such as Triton, and/or polyethylene glycol.

* * *

It is understood that the application of the teachings of the present invention to a specific problem or situation will be within the capabilities of one having ordinary skill in the art in light of the teachings contained herein. Examples of the products of the present invention appear below.

EXAMPLES

Materials and Methods

Reagents.

Recombinant soluble CD14 (rsCD14) and recombinant LBP (rLBP) were constructed and purified as

- 22 -

described (Hailman, E., et al. J. Exp. Med. **179**, 269-277 (1994)). Concentrations of all purified proteins were determined with a Micro BCA protein kit (Pierce, Rockford, IL) according to manufacturer's specification.

5 Since full-length rsCD14 terminates at position 348 of the mature protein (Hailman, E., et al. J. Exp. Med. **179**, 269-277 (1994)), we herein refer it as sCD14₁₋₃₄₈. The anti-CD14 mAb 3C10 was purified by chromatography on Protein G from the conditioned medium (CM) of a cell

10 line from American Type Culture Collection (ATCC TIB 228). Rabbit polyclonal anti-human CD14 antiserum was prepared as described (Juan, T. S. -C., et al. J. Biol. Chem. **270**, 1382-1387 (1995)). Rough LPS (*Salmonella* minnesota R60 or Re595) and smooth LPS (*E. coli* 0111:B4

15 or *Salmonella* minnesota wild-type) were purchased from LIST Biological Laboratories (Campbell, CA). Enzymes for DNA manipulation were purchased from Boehringer Mannheim (Indianapolis, IN).

20

Site-directed mutagenesis.

Nine alanine-substitution mutants of sCD14 were used in this study. FIG. 2 summarizes the names and the amino acid residues substituted in each mutant.

25 The Transformer site-directed mutagenesis kit (Clontech, Palo Alto, CA) was used as previously described (Juan, T. S. -C., et al. J. Biol. Chem. **270**, 5219-5224 (1995)) to generate cDNAs encoding alanine-substitution mutants of sCD14 cloned in a mammalian expression vector. The

30 primers used for each mutant are as follows:

5'-CGCCAGAACCTTGTGCAGCTGCCGCTGAAGATTCCGCTGC-3' for sCD14_{(7-10)A},

5'-GTGAGCTGGACGATGCAGCTGCCGCTGCGTCTGCAACTTC-3' for sCD14_{(11-14)A},

35 5'-CCGCTGCGTCTGCGCAGCTGCCGCACCTCAGCCGACTGG-3' for sCD14_{(18-21)A},

- 23 -

5'-GCAACTTCTCCGAAGCAGCTGCCGCCTGGTCCGAAGCCTTC-3' for
sCD14 (22-25)A,

5'-GAACCTCAGCCCGACGCAGCTGCAGCCTTCCAGTGTGTG-3' for
sCD14 (26-28)A,

5 5'-CCGACTGGTCCGAAGCAGCTGCGTGTGTGTCTGCAGTAGAG-3' for
sCD14 (30-31)A,

5'-CATGCCGGCGGTGCAGCTGCAGCGCCGTTTCTAAAGCGCG-3' for
sCD14 (45-48)A,

5'-GGTCTCAACCTAGAGGCAGCTGCAGCGCGCTCGATGCGGAC-3' for
10 sCD14 (49-52)A, and

5'-GAGCCGTTTCTAAAGGCAGCTGCTGCGGACGCCGACCCG-3' for
sCD14 (52-55)A.

15 Transient Expression of Mutant sCD14 Proteins in COS-7 Cells.

To express mutant sCD14 proteins, mammalian
expression vectors containing mutant sCD14 cDNAs were
introduced into COS-7 (ATCC CRL 1651) cells by
electroporation. Conditions for electroporation and
20 generation of serum-free CM from transfected COS-7 cells
were as described (Juan, T. S. -C., et al. J. Biol.
Chem. **270**, 1382-1387 (1995)). Expression of mutant
sCD14 was analyzed by Western blot using anti-CD14
polyclonal antibody.

25 BIAcore Analyses of Interactions Between sCD14 Mutants and 3C10 mAb.

Recognition of sCD14 mutant proteins by
neutralizing monoclonal antibody 3C10 was performed with
30 a BIAcore biosensor instrument. The instrument, CM5
sensor chips, and amine coupling kit were purchased from
Pharmacia Biosensor (Piscataway, NJ). Briefly, mAb 3C10
(200 µg/ml in 20 mM sodium acetate, pH 3.4) was
immobilized to a CM5 sensor chip by amine coupling
35 according to manufacturer's specifications. The flow
cell immobilized with 3C10 was then incubated in

- 24 -

succession with solutions as detailed in the following steps: Step 1, COS-7 CM for 2 min and Step 2, HBS buffer [10 mM N-2-hydroxyethylpiperazine-N'-2-ethanesulfonic acid, pH 7.5, 0.15 M NaCl, 3.4 mM EDTA, 0.005% (V/V) surfactant P20 (Pharmacia Biosensor)] for 2 min. For regeneration, 10 mM HCl solution was injected for 2 min. Injection was performed at a rate of 5 μ l/min. To quantitate the binding of sCD14 mutants in COS-7 CM to immobilized 3C10, we calculated relative a response unit (RRU). RRU was obtained by subtracting the response unit (RU) recorded just before injection of CM from the RU recorded after injection of CM and a 2 min wash.

Purification of sCD14₍₇₋₁₀₎A.

The expression vector containing the cDNA encoding sCD14₍₇₋₁₀₎A was stably transfected into Chinese hamster ovary (CHO) cells deficient in dihydrofolate reductase as described (Hailman, E., et al. J. Exp. Med. **179**, 269-277 (1994)). A single clone was grown without serum to generate CM containing sCD14₍₇₋₁₀₎A. Mutant protein was purified exactly as described (Juan, T. S. -C., et al. J. Biol. Chem. **270**, 5219-5224 (1995)) except immunoaffinity chromatography was performed with anti-CD14 polyclonal antibody coupled to Sepharose 4B (Pharmacia, Piscataway, NJ). Purity of the sample was checked by SDS-polyacrylamide gel electrophoresis (SDS-PAGE) followed by silver staining or Coomassie Blue staining. The changed amino acid sequence was verified through N-terminal sequencing.

U373 bioassays.

Growth of U373 cells (ATCC HTB17, Rockville, MD), activation by purified sCD14 preparations, and quantitation of IL-6 were performed exactly as described (Juan, T. S. -C., et al. J. Biol. Chem. **270**, 1382-1387 (1995)). Briefly, mixtures of sCD14₁₋₃₄₈ or sCD14₍₇₋₁₀₎A

- 25 -

and LPS were added to monolayers of U373 cells in serum--free medium and incubated for 24 h. IL-6 in the supernatant was then measured by ELISA.

5 Polymorphonuclear Leukocytes (PMN) Adhesion Assays.

The ability of rLBP and sCD14_{(7-10)A} or sCD14₁₋₃₄₈ to enable PMN adhesion to fibrinogen-coated plates was assessed by previously established protocols (Hailman, E., et al. J. Exp. Med. **179**, 269-277 (1994); Juan, T. S. -C., et al. J. Biol. Chem. **270**, 1382-1387 (1995)). Briefly, PMN were incubated for 10 min with LPS, rLBP, and sCD14_{(7-10)A} or sCD14₁₋₃₄₈, washed and adhesion to fibrinogen-coated surfaces was measured as described (Hailman, E., et al. J. Exp. Med. **179**, 269-277 (1994); Juan, T. S. -C., et al. J. Biol. Chem. **270**, 1382-1387 (1995)). When smooth LPS is used in this protocol, adhesion is completely dependent on addition of sCD14₁₋₃₄₈ (Juan, T. S. -C., et al. J. Biol. Chem. **270**, 1382-1387 (1995)).

The ability of sCD14_{(7-10)A} or sCD14₁₋₃₄₈ at high concentrations to bind LPS and inhibit LPS-mediated PMN adhesion was also assessed. In this experiment, rough LPS (*Salmonella minnesota* R60, 10 ng/ml) was incubated with rLBP (1 µg/ml) and the indicated concentrations of sCD14_{(7-10)A} or sCD14₁₋₃₄₈ for 30 min at 37°C before the addition of PMNs. The adhesion of PMNs was measured as described above.

30 Electrophoretic Mobility Shift Assays.

Whole cell extracts from U373 cells were prepared to assess transcription factor NF-κB activation. Cells were seeded in 6-well plates at a density of 1 million cells per well one day prior to stimulation. For stimulation, purified sCD14₁₋₃₄₈, sCD14_{Δ(57-64)} (Juan, T. S. -C., et al. J. Biol. Chem. **270**,

- 26 -

5219-5224 (1995)), or SCD14₍₇₋₁₀₎A was added at a final concentration of 20 ng/ml with or without 20 ng/ml of Re595 LPS for 20 h. Cells were washed twice with 1X PBS (GIBCO-BRL), and scraped in 200 µl of lysis buffer (20 mM HEPES, pH 7.9, 20% glycerol, 0.1 M KCl, 1 mM EDTA, 0.5 mM dithiothreitol, 1 mM Pefabloc (Boehringer Mannheim), 5 µg/ml Leupeptin, 1 mM sodium orthovanadate, and 2 µg/ml aprotinin) supplemented with 1% Triton X-100 (Sigma). Crude extracts were transferred to microfuge tubes and debris was separated by centrifugation at 14,000 X g for 10 min at 4°C. Extracts were quickly frozen in liquid nitrogen and stored at -80°C. Protein concentration of the whole cell extracts were determined by micro BCA assay and ranged between 1.5-2 µg/µl.

For examining the NF-κB complexes, we performed electrophoretic mobility shift assays. Two oligonucleotides:

5'-CATGGAGGGACTTTCGCTGGGGACTTTCAGC-3' and
5'-CATGGCTGGAAAGTCCCCAGCGAAAGTCCCTC-3'

were annealed to generate a double-stranded DNA containing the NF-κB binding site of human immunodeficient virus long terminal repeat promoter (Nabel, G. and Baltimore, D. Nature **326**, 711-713 (1987)). This annealed DNA fragment was then filled in with Klenow fragment (Boehringer Mannheim) and α-³²P dCTP (Amersham, Arlington Heights, IL) and used as probe at a concentration of 50,000 cpm per lane (about 25 fmole). For binding, 4 µl of whole cell extract was incubated with 4 µl of 5 X binding buffer (150 mM Tris-HCl, pH 8.0, 40 mM MgCl₂, 5 mM DTT, and 10% glycerol), 2.5 µg of (poly dI-dC):(poly dI-dC) (Pharmacia, Piscataway, NJ), radioactively labeled DNA probe and adequate amount of lysis buffer so that the final volume was 20 µl per reaction. The reactions were incubated in a 30°C water bath for 30 min and complexes were resolved in a native 4.5% polyacrylamide gel using 0.5 X TBE

- 27 -

(50 mM Tris-HCl, pH 8.0, 45 mM boric acid, and 5 mM EDTA) at 30 mA for 2 h. The gel was then vacuum-dried at 80°C for 1 h and exposed to Kodak X-ray film for 20 h. In competition experiments, 100 X molar excess of
5 unlabelled NF- κ B probe was pre-incubated for 10 min before addition of radioactive probe.

Native PAGE Assays.

To directly assess LPS-binding of purified
10 sCD14 preparations, sCD14₁₋₃₄₈ or sCD14_{(7-10)A} were incubated at various concentrations (0, 101, 303, and 909 nM) with 3 μ g/ml of ³H-LPS prepared from *E. coli* K12 strain LCD25 (List Biological Laboratories) in the presence or absence of 16.7 nM rLBP. The reaction was
15 incubated at 37°C for 30 min and then electrophoresed on native 4-20% polyacrylamide gels. Gels were prepared for fluorography as previously described (Hailman, E., et al. J. Exp. Med. **179**, 269-277 (1994)).

20 Inhibition of LPS-Induced TNF- α Production in Whole Blood.

The ability of sCD14 to bind LPS and inhibit TNF- α production in whole blood has been described (Haziot, A., et al. J. Immunol **152**, 5868-5876 (1994)).
25 Briefly, various concentrations of bovine serum albumin (Miles, New Haven, CT), sCD14₁₋₃₄₈, or sCD14_{(7-10)A} diluted in 50 μ l RPMI medium (GIBCO-BRL, Gaithersburg, MD) were added to 250 μ l of freshly-drawn blood using heparin as an anti-coagulant. Smooth LPS (*Salmonella minnesota*
30 wild-type) was added to a final concentration of 0.25 ng/ml. The reaction was incubated at 37°C for 3 h and supernatants were obtained by centrifugation at 16,000 Xg for 2 min. TNF- α concentrations in the supernatants were assayed using a Quantikine TNF- α ELISA kit (R & D
35 Systems, Minneapolis, MN) as suggested by the manufacturer.

- 28 -

EXAMPLE 1

Alanine Substitution at amino acids 7 to 10 or 11
5 to 14 Disrupts Binding of Neutralizing mAb 3C10
to CD14.

3C10 is a mAb that recognizes the N-terminal
152 amino acid of CD14 (Juan, T. S. -C., et al. J. Biol.
Chem. 270, 1382-1387 (1995)). Previous experiments have
10 shown that 3C10 neutralizes the activity of sCD14₁₋₃₄₈
(Wright, S. D., et al. Science 249, 1431-1433 (1990);
Hailman, E., et al. J. Exp. Med. 179, 269-277 (1994);
Frey, E. A., et al. J. Exp. Med. 176, 1665-1671 (1992);
Wright, S. D., et al. J. Exp. Med. 173, 1281-1286
15 (1991)). To verify that neutralization of sCD14
activity was due to binding of epitopes within the
N-terminal 152 amino acids, we demonstrated that 3C10
inhibited IL-6 production in U373 cells mediated by
either sCD14₁₋₃₄₈ or sCD14₁₋₁₅₂ (data not shown).

20 To map the epitope for mAb 3C10, a series of
alanine-substitution mutants were generated by site-
directed mutagenesis (FIG. 2). Plasmids containing cDNA
sequences encoding different sCD14 mutants were
transfected into COS-7 cells and CM from these cells
25 were examined for the expression of mutant sCD14
proteins by Western blot. With the exception of
sCD14_{(18-21)A}, all sCD14 mutants were expressed and
secreted by COS-7 cells (FIG. 2). BIAcore analysis
(FIG. 3) was then used to examine the ability of CM
30 containing mutant sCD14 to bind 3C10. CM containing
sCD14_{(7-10)A} or sCD14_{(11-14)A} were found not to bind 3C10.
These data suggest that the region between amino acids 7
and 14 is involved in recognizing 3C10.

35

- 29 -

EXAMPLE 2

Purification and characterization of sCD14₍₇₋₁₀₎A.

Since neutralizing mAb 3C10 recognized amino acids 7 to 14, we reasoned that this region of CD14 could play an important role in the biological activity of CD14. To help understand the role of this region, we generated a stable CHO cell line expressing sCD14₍₇₋₁₀₎A and purified mutant protein from the serum-free CM of this cell line. Purified sCD14₍₇₋₁₀₎A migrated with an apparent Mr of 55,000 when analyzed by reducing SDS-PAGE (data not shown). N-terminal sequencing confirmed that the amino acids between 7 and 10 were replaced with alanines residues.

15

EXAMPLE 3

mAb 3C10 Does not Recognize purified sCD14₍₇₋₁₀₎A.

BIACore realtime analysis was again used to determine whether mAb 3C10 is able to bind purified sCD14₍₇₋₁₀₎A. FIG. 4 shows that sCD14₁₋₃₄₈ recognized immobilized 3C10 and caused an increase of 1800 RU 2 min after wash (compare RU of the sensorgram before HCl injection at T=300 to that before injection of sCD14₁₋₃₄₈ at T=0), confirming previous observations (Juan, T. S. -C., et al. *J. Biol. Chem.* 270, 1382-1387 (1995)). However, purified sCD14₍₇₋₁₀₎A failed to recognize 3C10 and caused only slight RU change (compare RU of the sensorgram after second wash at T=750 to that before injection of sCD14₁₋₃₄₈ at T=0) similar to that observed when an irrelevant protein such as bovine serum albumin was injected (data not shown), demonstrating that amino acids 7-10 are required for mAb 3C10 binding.

35

- 30 -

EXAMPLE 4

sCD14₍₇₋₁₀₎A has Reduced Ability to Mediate Cellular Responses to LPS.

5 To assess the consequences of mutating residues between 7 and 10 in sCD14, we used two previously described assays (Hailman, E., et al. J. Exp. Med. **179**, 269-277 (1994); Juan, T. S. -C., et al. J. Biol. Chem. **270**, 1382-1387 (1995); Frey, E. A.,
10 et al. J. Exp. Med. **176**, 1665-1671 (1992)) to measure sCD14₍₇₋₁₀₎A bioactivity. We first examined the ability of sCD14₍₇₋₁₀₎A to enable responses of U373 cells to LPS. Addition of as little as 5 ng/ml sCD14₁₋₃₄₈ in the presence of LPS enabled strong IL-6 production
15 (FIG. 5A). In contrast, sCD14₍₇₋₁₀₎A was greatly impaired in its ability to enable responses, and required approximately 10-fold more protein in order to give a similar response to that of sCD14₁₋₃₄₈ (FIG. 5A).

We also examined whether sCD14₍₇₋₁₀₎A could enable
20 LPS-induced adhesion of PMN to fibrinogen. FIG. 5B shows that 100 ng/ml sCD14₁₋₃₄₈ enabled a strong adhesive response of PMN to smooth LPS and rLBP. However, very little response was seen even when 10,000 ng/ml sCD14₍₇₋₁₀₎A was added. These findings confirm that the
25 region between amino acids 7 and 10 are necessary for the biological activity of sCD14.

EXAMPLE 5

sCD14₍₇₋₁₀₎A is Impaired in its Ability to Activate Transcription Factor NF- κ B in the Presence of LPS.

LPS and sCD14-mediated activation of cells has been shown to involve activation of transcription
35 factors such as NF- κ B (Sen, R., and Baltimore, D. Cell **47**, 921-928 (1986); Lee, J. D., et al. J. Exp. Med.

- 31 -

175, 1697-1705 (1992); Bagasra, D., et al. Proc. Natl. Acad. Sci. U. S. A. **89**, 6285-6289 (1992)). To assess whether the mutation in sCD14_{(7-10)A} affected downstream signaling, we examined NF- κ B activation in U373 cells

5 treated with wildtype or mutant sCD14. In the absence of LPS or sCD14, U373 cells possess endogenous NF- κ B which forms a complex with labeled NF- κ B probe (Complex 1, FIG. 6, lane 1). Stimulation with LPS alone or sCD14₁₋₃₄₈ alone caused slight enhancement of NF- κ B

10 complex 1 and slight induction of a new NF- κ B complex (Complex 2, FIG. 6, lanes 2 and 3), but addition of sCD14₁₋₃₄₈ and LPS greatly induced both complexes of NF- κ B (FIG. 6, compare lanes 1 and 4). Both complexes 1 and 2 are NF- κ B specific since a 100-fold excess of unlabelled

15 NF- κ B oligonucleotide pre-incubated with extracts of U373 cells eliminated formation of both complexes (data not shown). Stimulation of U373 cells with sCD14_{(7-10)A} and LPS caused only 5% of NF- κ B activation as quantitated by gel scanning (FIG. 6, lane 6).

20 Comparatively, stimulation of U373 cells with a mutant which does not bind LPS (sCD14 Δ 57-64) [Note: Δ 57-64 means deletion of amino acids 57 - 64] failed to activate NF- κ B complexes even in the presence of LPS (FIG. 6, lane 8). These data indicate that a defect in

25 sCD14_{(7-10)A} is observed at the level at the transcription factor NF- κ B. Since activation of NF- κ B is an early event in signal transduction (Grilli, M., et al. Int. Rev. Cytol. **143**, 1-62 (1993)), these data suggest that sCD14_{(7-10)A} fails to enable signaling.

30

EXAMPLE 6

sCD14_{(7-10)A} Forms A Stable Complex with LPS.

Reduced signaling by sCD14_{(7-10)A} could be due

35 to a defect in binding LPS. To directly assess whether sCD14_{(7-10)A} binds LPS normally, we used a native PAGE

- 32 -

assay to detect stable complexes between sCD14₁₋₃₄₈ or sCD14_{(7-10)A} and ³H-LPS. As previously reported (Hailman, E., et al. J. Exp. Med. 179, 269-277 (1994)), formation of stable complexes between sCD14₁₋₃₄₈ and LPS could be
5 observed after 30 min of incubation (FIG. 7A), and addition of rLBP lowered the concentration of sCD14₁₋₃₄₈ required for complex formation (compare lane 2 of FIG. 7B to lane 2 of FIG. 7A). This is consistent with the previous observation (Hailman, E., et al. J. Exp. Med.
10 179, 269-277 (1994)) that rLBP accelerates the transfer of LPS to sCD14. Interestingly, sCD14_{(7-10)A} was also able to form stable complexes with ³H-LPS in the absence of rLBP (FIG 7A, lanes 5-7), and this complex formation was also facilitated by rLBP (compare lane 5 of FIG. 7B
15 to lane 5 of FIG. 7A). These data confirm that sCD14_{(7-10)A} is capable of binding LPS in an LBP-facilitated and in an LBP-independent fashion *in vitro* and suggest that the reduced biological activity of sCD14_{(7-10)A} is not due to an inability to bind LPS.

20

EXAMPLE 7

Inhibition of LPS-induced Cellular Responses by High Concentrations of sCD14.

25 To further confirm that sCD14_{(7-10)A} could bind LPS, we utilized two cell-based assays in which high concentrations of sCD14 prevent LPS-mediated activation of cells. In the first assay, sCD14₁₋₃₄₈ or sCD14_{(7-10)A} were tested for their ability to inhibit adhesion of PMN
30 to fibrinogen induced by LPS (FIG. 8A). In this experiment, constant concentrations of LPS and rLBP were incubated with increasing amounts (from 1 to 100 µg/ml) of sCD14₁₋₃₄₈ or sCD14_{(7-10)A}. Both proteins were capable of neutralizing LPS and inhibiting the adhesion of PMN
35 induced by LPS.

- 33 -

We also examined whether sCD14₍₇₋₁₀₎A could inhibit LPS-mediated TNF- α production in a whole blood assay, as has been shown for a recombinant sCD14 expressed in Baculovirus (Haziot, A., et al. J. Immunol **152**, 5868-5876 (1994)). Addition of increasing amounts of sCD14₁₋₃₄₈ or sCD14₍₇₋₁₀₎A caused inhibition of TNF- α production in the whole blood assay (FIG. 8B), while addition of bovine serum albumin did not inhibit TNF- α production, confirming the previous observation (Haziot, A., et al. J. Immunol **152**, 5868-5876 (1994)). These data confirm that sCD14₍₇₋₁₀₎A interacts with LPS as well as sCD14₁₋₃₄₈.

Discussion of Examples 1 - 7

In the above examples, we mapped the epitope for neutralizing mAb 3C10 to the region between amino acids 7 and 14 of sCD14. Substitution of alanine residues in this region prevented binding of 3C10 to sCD14. These data are consistent with our previous finding (Juan, T. S. -C., et al. J. Biol. Chem. **270**, 1382-1387 (1995); Juan, T. S. -C., et al. J. Biol. Chem. **270**, 5219-5224 (1995)) that the 3C10 epitope is located within the first 152 amino acids of sCD14 and is distinct from the epitope of MEM-18 at residues 57-64. To help understand how the 3C10 epitope contributes to CD14 function, we purified sCD14₍₇₋₁₀₎A and showed that this protein was severely impaired in its ability to activate cells. Inability of this protein to promote activation of NF- κ B suggests that sCD14₍₇₋₁₀₎A fails to support LPS-mediated signaling.

The defect in sCD14₍₇₋₁₀₎A signaling is unlikely to result from an inability of this protein to bind LPS properly or to interact with LBP. sCD14₍₇₋₁₀₎A binds LPS normally, as examined by gelshift (FIG. 7A) and two cell-based assays (FIG. 8) and rLBP facilitates transfer

- 34 -

of LPS to sCD14_{(7-10)A} (FIG. 7B). These data confirm our previous observation that 3C10 binds normally to complexes of sCD14 and LPS (Juan, T. S. -C., et al. J. Biol. Chem. 270, 5219-5224 (1995)). These experiments
5 measured direct binding of LPS to sCD14_{(7-10)A}, not the binding of LPS-LBP complexes to cell surface CD14 measured in other reports (Wright, S. D., et al. Science 249: 1431-1433 (1990); Viriyakosol, S. and Kirkland, T. N. J. Biol. Chem. 270, 361-368 (1995)).
10 Since sCD14_{(7-10)A} binds LPS normally, its defect in signaling is likely to be manifest at the cell membrane. We (Frey, E. A., et al. J. Exp. Med. 176, 1665-1671 (1992)) and others (Pugin, J., et al. Proc. Natl. Acad. Sci. U. S. A. 90, 2744-2748 (1993); Haziot, A., et al. J. Immunol. 151, 1500-1507 (1993); Arditi, M., et al. Infect. Immun. 61, 3149-3156 (1993)) have
15 postulated the existence of a transmembrane protein that interacts with LPS and/or CD14 and transmits signals to the cytoplasm. It is thus possible that residues 7-10
20 are essential for the interaction of sCD14 with this transmembrane constituent. Alternatively, sCD14_{(7-10)A} may be defective in delivering LPS to the lipid bilayer of cells. We have recently shown sCD14 rapidly shuttles LPS into HDL particles (Wurfel, M. M., et al. J. Exp. Med. 181: 1743-1754 (1995)) and into
25 phospholipid vesicles (M. M. Wurfel and S. D. W., manuscript in preparation), and it is thus possible that residues 7-10 are essential for delivery of bound LPS into the plasma membrane of cells.

30

EXAMPLE 8

Gram Positive Cell Components Compete with LPS for Binding to sCD14

35 FIG. 9 presents the evidence that a gram-positive molecule present in the phenol extract of S.

- 35 -

aureus (SACE) can bind to sCD14 and compete with LPS for a binding site. Other data (not shown) indicates that SACE strongly stimulates cells in a CD14-dependent fashion. The binding site(s) now defined on CD14 may be relevant not only to responses initiated by gram-negative but also by gram-positive bacteria.

Abbreviations

The abbreviations used in the Examples section above are: BCIP, 5-bromo-4-chloro-3-indoyl phosphate-toluidine salt; BPI, bactericidal/permeability-increasing protein; CHO, Chinese hamster ovary; CD, circular dichroism; CM, conditioned medium; HBSS, Hank's balanced salt solution; IL-6, interleukin-6; LALF, *Limulus* anti-LPS factor; LBP, LPS-binding protein; LPS, lipopolysaccharide; NBT, p-nitro blue tetrazolium chloride; PAGE, polyacrylamide gel electrophoresis; PBS, phosphate-buffered saline; PMN, polymorphonuclear leukocyte; r, recombinant; RU, response unit; sCD14, soluble CD14; ELISA, enzyme linked immunosorbant assay.

* * *

The invention now being fully described, it will be apparent to one of ordinary skill in the art that many changes and modifications can be made thereto, without departing from the spirit and scope of the invention as set forth herein.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

- (i) APPLICANT: Juan, Shao-Chieh
Lichenstein, Henri S.
Wright, Samuel D.
- (ii) TITLE OF INVENTION: ANTI-INFLAMMATORY CD14
PEPTIDES
- (iii) NUMBER OF SEQUENCES: 37
- (iv) CORRESPONDENCE ADDRESS:
 - (A) ADDRESSEE: Amgen Inc.
 - (B) STREET: 1840 S. Dehavilland Drive
 - (C) CITY: Thousand Oaks
 - (D) STATE: CA
 - (E) COUNTRY: USA
 - (F) ZIP: 91320-1789
- (v) COMPUTER READABLE FORM:
 - (A) MEDIUM TYPE: Floppy disk
 - (B) COMPUTER: IBM PC compatible
 - (C) OPERATING SYSTEM: PC-DOS/MS-DOS
 - (D) SOFTWARE: PatentIn Release #1.0, Version
#1.25
- (vi) CURRENT APPLICATION DATA:
 - (A) APPLICATION NUMBER: US 08/484,397
 - (B) FILING DATE: 07-JUN-1995
 - (C) CLASSIFICATION:
- (viii) ATTORNEY/AGENT INFORMATION:
 - (A) NAME: Cook Ph.D., Robert R.
 - (B) REGISTRATION NUMBER: 31,602
 - (C) REFERENCE/DOCKET NUMBER: A-324A
- (ix) TELECOMMUNICATION INFORMATION:
 - (A) TELEPHONE: 805 447-1000
 - (B) TELEFAX: 805/499-8011

(2) INFORMATION FOR SEQ ID NO:1:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 4 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

Glu Leu Asp Asp
1

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 348 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Thr	Thr	Pro	Glu	Pro	Cys	Gly	Gly	Gly	Gly	Glu	Asp	Phe	Arg	Cys	Val	1	5	10	15
Cys	Asn	Phe	Ser	Glu	Pro	Gln	Pro	Asp	Trp	Ser	Glu	Ala	Phe	Gln	Cys	20	25	30	
Val	Ser	Ala	Val	Glu	Val	Glu	Ile	His	Ala	Gly	Gly	Leu	Asn	Leu	Glu	35	40	45	
Pro	Phe	Leu	Lys	Arg	Val	Asp	Ala	Asp	Ala	Asp	Pro	Arg	Gln	Tyr	Ala	50	55	60	
Asp	Thr	Val	Lys	Ala	Leu	Arg	Val	Arg	Arg	Leu	Thr	Val	Gly	Ala	Ala	65	70	75	80
Gln	Val	Pro	Ala	Gln	Leu	Leu	Val	Gly	Ala	Leu	Arg	Val	Leu	Ala	Tyr	85	90	95	
Ser	Arg	Leu	Lys	Glu	Leu	Thr	Leu	Glu	Asp	Leu	Lys	Ile	Thr	Gly	Thr	100	105	110	
Met	Pro	Pro	Leu	Pro	Leu	Glu	Ala	Thr	Gly	Leu	Ala	Leu	Ser	Ser	Leu	115	120	125	
Arg	Leu	Arg	Asn	Val	Ser	Trp	Ala	Thr	Gly	Arg	Ser	Trp	Leu	Ala	Glu	130	135	140	
Leu	Gln	Gln	Trp	Leu	Lys	Pro	Gly	Leu	Lys	Val	Leu	Ser	Ile	Ala	Gln	145	150	155	160

Ala His Ser Pro Ala Phe Ser Cys Glu Gln Val Arg Ala Phe Pro Ala
 165 170 175
 Leu Thr Ser Leu Asp Leu Ser Asp Asn Pro Gly Leu Gly Glu Arg Gly
 180 185 190
 Leu Met Ala Ala Leu Cys Pro His Lys Phe Pro Ala Ile Gln Asn Leu
 195 200 205
 Ala Leu Arg Asn Thr Gly Met Glu Thr Pro Thr Gly Val Cys Ala Ala
 210 215 220
 Leu Ala Ala Ala Gly Val Gln Pro His Ser Leu Asp Leu Ser His Asn
 225 230 235 240
 Ser Leu Arg Ala Thr Val Asn Pro Ser Ala Pro Arg Cys Met Trp Ser
 245 250 255
 Ser Ala Leu Asn Ser Leu Asn Leu Ser Phe Ala Gly Leu Glu Gln Val
 260 265 270
 Pro Lys Gly Leu Pro Ala Lys Leu Arg Val Leu Asp Leu Ser Cys Asn
 275 280 285
 Arg Leu Asn Arg Ala Pro Gln Pro Asp Glu Leu Pro Glu Val Asp Asn
 290 295 300
 Leu Thr Leu Asp Gly Asn Pro Phe Leu Val Pro Gly Thr Ala Leu Pro
 305 310 315 320
 His Glu Gly Ser Met Asn Ser Gly Val Val Pro Ala Cys Ala Arg Ser
 325 330 335
 Thr Leu Ser Val Gly Val Ser Gly Thr Leu Val Leu
 340 345

(2) INFORMATION FOR SEQ ID NO:3:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 348 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

Thr Thr Pro Glu Pro Cys Ala Ala Ala Ala Glu Asp Phe Arg Cys Val
 1 5 10 15

Cys Asn Phe Ser Glu Pro Gln Pro Asp Trp Ser Glu Ala Phe Gln Cys
 20 25 30
 Val Ser Ala Val Glu Val Glu Ile His Ala Gly Gly Leu Asn Leu Glu
 35 40 45
 Pro Phe Leu Lys Arg Val Asp Ala Asp Ala Asp Pro Arg Gln Tyr Ala
 50 55 60
 Asp Thr Val Lys Ala Leu Arg Val Arg Arg Leu Thr Val Gly Ala Ala
 65 70 75 80
 Gln Val Pro Ala Gln Leu Leu Val Gly Ala Leu Arg Val Leu Ala Tyr
 85 90 95
 Ser Arg Leu Lys Glu Leu Thr Leu Glu Asp Leu Lys Ile Thr Gly Thr
 100 105 110
 Met Pro Pro Leu Pro Leu Glu Ala Thr Gly Leu Ala Leu Ser Ser Leu
 115 120 125
 Arg Leu Arg Asn Val Ser Trp Ala Thr Gly Arg Ser Trp Leu Ala Glu
 130 135 140
 Leu Gln Gln Trp Leu Lys Pro Gly Leu Lys Val Leu Ser Ile Ala Gln
 145 150 155 160
 Ala His Ser Pro Ala Phe Ser Cys Glu Gln Val Arg Ala Phe Pro Ala
 165 170 175
 Leu Thr Ser Leu Asp Leu Ser Asp Asn Pro Gly Leu Gly Glu Arg Gly
 180 185 190
 Leu Met Ala Ala Leu Cys Pro His Lys Phe Pro Ala Ile Gln Asn Leu
 195 200 205
 Ala Leu Arg Asn Thr Gly Met Glu Thr Pro Thr Gly Val Cys Ala Ala
 210 215 220
 Leu Ala Ala Ala Gly Val Gln Pro His Ser Leu Asp Leu Ser His Asn
 225 230 235 240
 Ser Leu Arg Ala Thr Val Asn Pro Ser Ala Pro Arg Cys Met Trp Ser
 245 250 255
 Ser Ala Leu Asn Ser Leu Asn Leu Ser Phe Ala Gly Leu Glu Gln Val
 260 265 270
 Pro Lys Gly Leu Pro Ala Lys Leu Arg Val Leu Asp Leu Ser Cys Asn
 275 280 285
 Arg Leu Asn Arg Ala Pro Gln Pro Asp Glu Leu Pro Glu Val Asp Asn
 290 295 300

Leu Thr Leu Asp Gly Asn Pro Phe Leu Val Pro Gly Thr Ala Leu Pro
 305 310 315 320
 His Glu Gly Ser Met Asn Ser Gly Val Val Pro Ala Cys Ala Arg Ser
 325 330 335
 Thr Leu Ser Val Gly Val Ser Gly Thr Leu Val Leu
 340 345

(2) INFORMATION FOR SEQ ID NO:4:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 348 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

Thr Thr Pro Glu Pro Cys Val Val Val Val Glu Asp Phe Arg Cys Val
 1 5 10
 Cys Asn Phe Ser Glu Pro Gln Pro Asp Trp Ser Glu Ala Phe Gln Cys
 20 25 30
 Val Ser Ala Val Glu Val Glu Ile His Ala Gly Gly Leu Asn Leu Glu
 35 40 45
 Pro Phe Leu Lys Arg Val Asp Ala Asp Ala Asp Pro Arg Gln Tyr Ala
 50 55 60
 Asp Thr Val Lys Ala Leu Arg Val Arg Arg Leu Thr Val Gly Ala Ala
 65 70 75 80
 Gln Val Pro Ala Gln Leu Leu Val Gly Ala Leu Arg Val Leu Ala Tyr
 85 90 95
 Ser Arg Leu Lys Glu Leu Thr Leu Glu Asp Leu Lys Ile Thr Gly Thr
 100 105 110
 Met Pro Pro Leu Pro Leu Glu Ala Thr Gly Leu Ala Leu Ser Ser Leu
 115 120 125
 Arg Leu Arg Asn Val Ser Trp Ala Thr Gly Arg Ser Trp Leu Ala Glu
 130 135 140
 Leu Gln Gln Trp Leu Lys Pro Gly Leu Lys Val Leu Ser Ile Ala Gln
 145 150 155 160

Ala His Ser Pro Ala Phe Ser Cys Glu Gln Val Arg Ala Phe Pro Ala
 165 170 175
 Leu Thr Ser Leu Asp Leu Ser Asp Asn Pro Gly Leu Gly Glu Arg Gly
 180 185 190
 Leu Met Ala Ala Leu Cys Pro His Lys Phe Pro Ala Ile Gln Asn Leu
 195 200 205
 Ala Leu Arg Asn Thr Gly Met Glu Thr Pro Thr Gly Val Cys Ala Ala
 210 215 220
 Leu Ala Ala Ala Gly Val Gln Pro His Ser Leu Asp Leu Ser His Asn
 225 230 235 240
 Ser Leu Arg Ala Thr Val Asn Pro Ser Ala Pro Arg Cys Met Trp Ser
 245 250 255
 Ser Ala Leu Asn Ser Leu Asn Leu Ser Phe Ala Gly Leu Glu Gln Val
 260 265 270
 Pro Lys Gly Leu Pro Ala Lys Leu Arg Val Leu Asp Leu Ser Cys Asn
 275 280 285
 Arg Leu Asn Arg Ala Pro Gln Pro Asp Glu Leu Pro Glu Val Asp Asn
 290 295 300
 Leu Thr Leu Asp Gly Asn Pro Phe Leu Val Pro Gly Thr Ala Leu Pro
 305 310 315 320
 His Glu Gly Ser Met Asn Ser Gly Val Val Pro Ala Cys Ala Arg Ser
 325 330 335
 Thr Leu Ser Val Gly Val Ser Gly Thr Leu Val Leu
 340 345

(2) INFORMATION FOR SEQ ID NO:5:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 348 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

Thr Thr Pro Glu Pro Cys Leu Leu Leu Leu Glu Asp Phe Arg Cys Val
 1 5 10 15

Cys Asn Phe Ser Glu Pro Gln Pro Asp Trp Ser Glu Ala Phe Gln Cys
 20 25 30
 Val Ser Ala Val Glu Val Glu Ile His Ala Gly Gly Leu Asn Leu Glu
 35 40 45
 Pro Phe Leu Lys Arg Val Asp Ala Asp Ala Asp Pro Arg Gln Tyr Ala
 50 55 60
 Asp Thr Val Lys Ala Leu Arg Val Arg Arg Leu Thr Val Gly Ala Ala
 65 70 75 80
 Gln Val Pro Ala Gln Leu Leu Val Gly Ala Leu Arg Val Leu Ala Tyr
 85 90 95
 Ser Arg Leu Lys Glu Leu Thr Leu Glu Asp Leu Lys Ile Thr Gly Thr
 100 105 110
 Met Pro Pro Leu Pro Leu Glu Ala Thr Gly Leu Ala Leu Ser Ser Leu
 115 120 125
 Arg Leu Arg Asn Val Ser Trp Ala Thr Gly Arg Ser Trp Leu Ala Glu
 130 135 140
 Leu Gln Gln Trp Leu Lys Pro Gly Leu Lys Val Leu Ser Ile Ala Gln
 145 150 155 160
 Ala His Ser Pro Ala Phe Ser Cys Glu Gln Val Arg Ala Phe Pro Ala
 165 170 175
 Leu Thr Ser Leu Asp Leu Ser Asp Asn Pro Gly Leu Gly Glu Arg Gly
 180 185 190
 Leu Met Ala Ala Leu Cys Pro His Lys Phe Pro Ala Ile Gln Asn Leu
 195 200 205
 Ala Leu Arg Asn Thr Gly Met Glu Thr Pro Thr Gly Val Cys Ala Ala
 210 215 220
 Leu Ala Ala Ala Gly Val Gln Pro His Ser Leu Asp Leu Ser His Asn
 225 230 235 240
 Ser Leu Arg Ala Thr Val Asn Pro Ser Ala Pro Arg Cys Met Trp Ser
 245 250 255
 Ser Ala Leu Asn Ser Leu Asn Leu Ser Phe Ala Gly Leu Glu Gln Val
 260 265 270
 Pro Lys Gly Leu Pro Ala Lys Leu Arg Val Leu Asp Leu Ser Cys Asn
 275 280 285
 Arg Leu Asn Arg Ala Pro Gln Pro Asp Glu Leu Pro Glu Val Asp Asn

290 295 300
 Leu Thr Leu Asp Gly Asn Pro Phe Leu Val Pro Gly Thr Ala Leu Pro
 305 310 315 320
 His Glu Gly Ser Met Asn Ser Gly Val Val Pro Ala Cys Ala Arg Ser
 325 330 335
 Thr Leu Ser Val Gly Val Ser Gly Thr Leu Val Leu
 340 345

(2) INFORMATION FOR SEQ ID NO:6:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 348 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

Thr Thr Pro Glu Pro Cys Ile Ile Ile Ile Glu Asp Phe Arg Cys Val
 1 5 10
 Cys Asn Phe Ser Glu Pro Gln Pro Asp Trp Ser Glu Ala Phe Gln Cys
 20 25 30
 Val Ser Ala Val Glu Val Glu Ile His Ala Gly Gly Leu Asn Leu Glu
 35 40 45
 Pro Phe Leu Lys Arg Val Asp Ala Asp Ala Asp Pro Arg Gln Tyr Ala
 50 55 60
 Asp Thr Val Lys Ala Leu Arg Val Arg Arg Leu Thr Val Gly Ala Ala
 65 70 75 80
 Gln Val Pro Ala Gln Leu Leu Val Gly Ala Leu Arg Val Leu Ala Tyr
 85 90 95
 Ser Arg Leu Lys Glu Leu Thr Leu Glu Asp Leu Lys Ile Thr Gly Thr
 100 105 110
 Met Pro Pro Leu Pro Leu Glu Ala Thr Gly Leu Ala Leu Ser Ser Leu
 115 120 125
 Arg Leu Arg Asn Val Ser Trp Ala Thr Gly Arg Ser Trp Leu Ala Glu
 130 135 140
 Leu Gln Gln Trp Leu Lys Pro Gly Leu Lys Val Leu Ser Ile Ala Gln
 145 150 155 160

- 44 -

Ala His Ser Pro Ala Phe Ser Cys Glu Gln Val Arg Ala Phe Pro Ala
 165 170 175
 Leu Thr Ser Leu Asp Leu Ser Asp Asn Pro Gly Leu Gly Glu Arg Gly
 180 185 190
 Leu Met Ala Ala Leu Cys Pro His Lys Phe Pro Ala Ile Gln Asn Leu
 195 200 205
 Ala Leu Arg Asn Thr Gly Met Glu Thr Pro Thr Gly Val Cys Ala Ala
 210 215 220
 Leu Ala Ala Ala Gly Val Gln Pro His Ser Leu Asp Leu Ser His Asn
 225 230 235 240
 Ser Leu Arg Ala Thr Val Asn Pro Ser Ala Pro Arg Cys Met Trp Ser
 245 250 255
 Ser Ala Leu Asn Ser Leu Asn Leu Ser Phe Ala Gly Leu Glu Gln Val
 260 265 270
 Pro Lys Gly Leu Pro Ala Lys Leu Arg Val Leu Asp Leu Ser Cys Asn
 275 280 285
 Arg Leu Asn Arg Ala Pro Gln Pro Asp Glu Leu Pro Glu Val Asp Asn
 290 295 300
 Leu Thr Leu Asp Gly Asn Pro Phe Leu Val Pro Gly Thr Ala Leu Pro
 305 310 315 320
 His Glu Gly Ser Met Asn Ser Gly Val Val Pro Ala Cys Ala Arg Ser
 325 330 335
 Thr Leu Ser Val Gly Val Ser Gly Thr Leu Val Leu
 340 345

(2) INFORMATION FOR SEQ ID NO:7:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 348 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

Thr Thr Pro Glu Pro Cys Pro Pro Pro Pro Glu Asp Phe Arg Cys Val
 1 5 10 15

Cys Asn Phe Ser Glu Pro Gln Pro Asp Trp Ser Glu Ala Phe Gln Cys
 20 25 30
 Val Ser Ala Val Glu Val Glu Ile His Ala Gly Gly Leu Asn Leu Glu
 35 40 45
 Pro Phe Leu Lys Arg Val Asp Ala Asp Ala Asp Pro Arg Gln Tyr Ala
 50 55 60
 Asp Thr Val Lys Ala Leu Arg Val Arg Arg Leu Thr Val Gly Ala Ala
 65 70 75 80
 Gln Val Pro Ala Gln Leu Leu Val Gly Ala Leu Arg Val Leu Ala Tyr
 85 90 95
 Ser Arg Leu Lys Glu Leu Thr Leu Glu Asp Leu Lys Ile Thr Gly Thr
 100 105 110
 Met Pro Pro Leu Pro Leu Glu Ala Thr Gly Leu Ala Leu Ser Ser Leu
 115 120 125
 Arg Leu Arg Asn Val Ser Trp Ala Thr Gly Arg Ser Trp Leu Ala Glu
 130 135 140
 Leu Gln Gln Trp Leu Lys Pro Gly Leu Lys Val Leu Ser Ile Ala Gln
 145 150 155 160
 Ala His Ser Pro Ala Phe Ser Cys Glu Gln Val Arg Ala Phe Pro Ala
 165 170 175
 Leu Thr Ser Leu Asp Leu Ser Asp Asn Pro Gly Leu Gly Glu Arg Gly
 180 185 190
 Leu Met Ala Ala Leu Cys Pro His Lys Phe Pro Ala Ile Gln Asn Leu
 195 200 205
 Ala Leu Arg Asn Thr Gly Met Glu Thr Pro Thr Gly Val Cys Ala Ala
 210 215 220
 Leu Ala Ala Ala Gly Val Gln Pro His Ser Leu Asp Leu Ser His Asn
 225 230 235 240
 Ser Leu Arg Ala Thr Val Asn Pro Ser Ala Pro Arg Cys Met Trp Ser
 245 250 255
 Ser Ala Leu Asn Ser Leu Asn Leu Ser Phe Ala Gly Leu Glu Gln Val
 260 265 270
 Pro Lys Gly Leu Pro Ala Lys Leu Arg Val Leu Asp Leu Ser Cys Asn
 275 280 285
 Arg Leu Asn Arg Ala Pro Gln Pro Asp Glu Leu Pro Glu Val Asp Asn
 290 295 300

- 46 -

Leu Thr Leu Asp Gly Asn Pro Phe Leu Val Pro Gly Thr Ala Leu Pro
 305 310 315 320

His Glu Gly Ser Met Asn Ser Gly Val Val Pro Ala Cys Ala Arg Ser
 325 330 335

Thr Leu Ser Val Gly Val Ser Gly Thr Leu Val Leu
 340 345

(2) INFORMATION FOR SEQ ID NO:8:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 334 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

Cys Val Cys Asn Phe Ser Glu Pro Gln Pro Asp Trp Ser Glu Ala Phe
 1 5 10 15

Gln Cys Val Ser Ala Val Glu Val Glu Ile His Ala Gly Gly Leu Asn
 20 25 30

Leu Glu Pro Phe Leu Lys Arg Val Asp Ala Asp Ala Asp Pro Arg Gln
 35 40 45

Tyr Ala Asp Thr Val Lys Ala Leu Arg Val Arg Arg Leu Thr Val Gly
 50 55 60

Ala Ala Gln Val Pro Ala Gln Leu Leu Val Gly Ala Leu Arg Val Leu
 65 70 75 80

Ala Tyr Ser Arg Leu Lys Glu Leu Thr Leu Glu Asp Leu Lys Ile Thr
 85 90 95

Gly Thr Met Pro Pro Leu Pro Leu Glu Ala Thr Gly Leu Ala Leu Ser
 100 105 110

Ser Leu Arg Leu Arg Asn Val Ser Trp Ala Thr Gly Arg Ser Trp Leu
 115 120 125

Ala Glu Leu Gln Gln Trp Leu Lys Pro Gly Leu Lys Val Leu Ser Ile
 130 135 140

Ala Gln Ala His Ser Pro Ala Phe Ser Cys Glu Gln Val Arg Ala Phe
 145 150 155 160

Pro Ala Leu Thr Ser Leu Asp Leu Ser Asp Asn Pro Gly Leu Gly Glu
 165 170 175
 Arg Gly Leu Met Ala Ala Leu Cys Pro His Lys Phe Pro Ala Ile Gln
 180 185 190
 Asn Leu Ala Leu Arg Asn Thr Gly Met Glu Thr Pro Thr Gly Val Cys
 195 200 205
 Ala Ala Leu Ala Ala Ala Gly Val Gln Pro His Ser Leu Asp Leu Ser
 210 215 220
 His Asn Ser Leu Arg Ala Thr Val Asn Pro Ser Ala Pro Arg Cys Met
 225 230 235 240
 Trp Ser Ser Ala Leu Asn Ser Leu Asn Leu Ser Phe Ala Gly Leu Glu
 245 250 255
 Gln Val Pro Lys Gly Leu Pro Ala Lys Leu Arg Val Leu Asp Leu Ser
 260 265 270
 Cys Asn Arg Leu Asn Arg Ala Pro Gln Pro Asp Glu Leu Pro Glu Val
 275 280 285
 Asp Asn Leu Thr Leu Asp Gly Asn Pro Phe Leu Val Pro Gly Thr Ala
 290 295 300
 Leu Pro His Glu Gly Ser Met Asn Ser Gly Val Val Pro Ala Cys Ala
 305 310 315 320
 Arg Ser Thr Leu Ser Val Gly Val Ser Gly Thr Leu Val Leu
 325 330

(2) INFORMATION FOR SEQ ID NO:9:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 152 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

Thr Thr Pro Glu Pro Cys Gly Gly Gly Glu Asp Phe Arg Cys Val
 1 5 10
 Cys Asn Phe Ser Glu Pro Gln Pro Asp Trp Ser Glu Ala Phe Gln Cys
 20 25 30

Val Ser Ala Val Glu Val Glu Ile His Ala Gly Gly Leu Asn Leu Glu
 35 40 45
 Pro Phe Leu Lys Arg Val Asp Ala Asp Ala Asp Pro Arg Gln Tyr Ala
 50 55 60
 Asp Thr Val Lys Ala Leu Arg Val Arg Arg Leu Thr Val Gly Ala Ala
 65 70 75 80
 Gln Val Pro Ala Gln Leu Leu Val Gly Ala Leu Arg Val Leu Ala Tyr
 85 90 95
 Ser Arg Leu Lys Glu Leu Thr Leu Glu Asp Leu Lys Ile Thr Gly Thr
 100 105 110
 Met Pro Pro Leu Pro Leu Glu Ala Thr Gly Leu Ala Leu Ser Ser Leu
 115 120 125
 Arg Leu Arg Asn Val Ser Trp Ala Thr Gly Arg Ser Trp Leu Ala Glu
 130 135 140
 Leu Gln Gln Trp Leu Lys Pro Gly
 145 150

(2) INFORMATION FOR SEQ ID NO:10:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 152 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

Thr Thr Pro Glu Pro Cys Ala Ala Ala Glu Asp Phe Arg Cys Val
 1 5 10 15
 Cys Asn Phe Ser Glu Pro Gln Pro Asp Trp Ser Glu Ala Phe Gln Cys
 20 25 30
 Val Ser Ala Val Glu Val Glu Ile His Ala Gly Gly Leu Asn Leu Glu
 35 40 45
 Pro Phe Leu Lys Arg Val Asp Ala Asp Ala Asp Pro Arg Gln Tyr Ala
 50 55 60
 Asp Thr Val Lys Ala Leu Arg Val Arg Arg Leu Thr Val Gly Ala Ala
 65 70 75 80

Gln Val Pro Ala Gln Leu Leu Val Gly Ala Leu Arg Val Leu Ala Tyr
 85 90 95

Ser Arg Leu Lys Glu Leu Thr Leu Glu Asp Leu Lys Ile Thr Gly Thr
 100 105 110

Met Pro Pro Leu Pro Leu Glu Ala Thr Gly Leu Ala Leu Ser Ser Leu
 115 120 125

Arg Leu Arg Asn Val Ser Trp Ala Thr Gly Arg Ser Trp Leu Ala Glu
 130 135 140

Leu Gln Gln Trp Leu Lys Pro Gly
 145 150

(2) INFORMATION FOR SEQ ID NO:11:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 152 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

Thr Thr Pro Glu Pro Cys Val Val Val Val Glu Asp Phe Arg Cys Val
 1 5 10 15

Cys Asn Phe Ser Glu Pro Gln Pro Asp Trp Ser Glu Ala Phe Gln Cys
 20 25 30

Val Ser Ala Val Glu Val Glu Ile His Ala Gly Gly Leu Asn Leu Glu
 35 40 45

Pro Phe Leu Lys Arg Val Asp Ala Asp Ala Asp Pro Arg Gln Tyr Ala
 50 55 60

Asp Thr Val Lys Ala Leu Arg Val Arg Arg Leu Thr Val Gly Ala Ala
 65 70 75 80

Gln Val Pro Ala Gln Leu Leu Val Gly Ala Leu Arg Val Leu Ala Tyr
 85 90 95

Ser Arg Leu Lys Glu Leu Thr Leu Glu Asp Leu Lys Ile Thr Gly Thr
 100 105 110

Met Pro Pro Leu Pro Leu Glu Ala Thr Gly Leu Ala Leu Ser Ser Leu
 115 120 125

- 50 -

Arg Leu Arg Asn Val Ser Trp Ala Thr Gly Arg Ser Trp Leu Ala Glu
130 135 140

Leu Gln Gln Trp Leu Lys Pro Gly
145 150

(2) INFORMATION FOR SEQ ID NO:12:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 152 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

Thr Thr Pro Glu Pro Cys Leu Leu Leu Leu Glu Asp Phe Arg Cys Val
1 5 10
Cys Asn Phe Ser Glu Pro Gln Pro Asp Trp Ser Glu Ala Phe Gln Cys
20 25 30
Val Ser Ala Val Glu Val Glu Ile His Ala Gly Gly Leu Asn Leu Glu
35 40 45
Pro Phe Leu Lys Arg Val Asp Ala Asp Ala Asp Pro Arg Gln Tyr Ala
50 55 60
Asp Thr Val Lys Ala Leu Arg Val Arg Arg Leu Thr Val Gly Ala Ala
65 70 75 80
Gln Val Pro Ala Gln Leu Leu Val Gly Ala Leu Arg Val Leu Ala Tyr
85 90 95
Ser Arg Leu Lys Glu Leu Thr Leu Glu Asp Leu Lys Ile Thr Gly Thr
100 105 110
Met Pro Pro Leu Pro Leu Glu Ala Thr Gly Leu Ala Leu Ser Ser Leu
115 120 125
Arg Leu Arg Asn Val Ser Trp Ala Thr Gly Arg Ser Trp Leu Ala Glu
130 135 140
Leu Gln Gln Trp Leu Lys Pro Gly
145 150

- 51 -

(2) INFORMATION FOR SEQ ID NO:13:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 152 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

```

Thr Thr Pro Glu Pro Cys Ile Ile Ile Ile Glu Asp Phe Arg Cys Val
1          5          10          15
Cys Asn Phe Ser Glu Pro Gln Pro Asp Trp Ser Glu Ala Phe Gln Cys
20          25          30
Val Ser Ala Val Glu Val Glu Ile His Ala Gly Gly Leu Asn Leu Glu
35          40          45
Pro Phe Leu Lys Arg Val Asp Ala Asp Ala Asp Pro Arg Gln Tyr Ala
50          55          60
Asp Thr Val Lys Ala Leu Arg Val Arg Arg Leu Thr Val Gly Ala Ala
65          70          75          80
Gln Val Pro Ala Gln Leu Leu Val Gly Ala Leu Arg Val Leu Ala Tyr
85          90          95
Ser Arg Leu Lys Glu Leu Thr Leu Glu Asp Leu Lys Ile Thr Gly Thr
100          105          110
Met Pro Pro Leu Pro Leu Glu Ala Thr Gly Leu Ala Leu Ser Ser Leu
115          120          125
Arg Leu Arg Asn Val Ser Trp Ala Thr Gly Arg Ser Trp Leu Ala Glu
130          135          140
Leu Gln Gln Trp Leu Lys Pro Gly
145          150

```

(2) INFORMATION FOR SEQ ID NO:14:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 152 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

Thr Thr Pro Glu Pro Cys Pro Pro Pro Pro Glu Asp Phe Arg Cys Val
 1 5 10 15
 Cys Asn Phe Ser Glu Pro Gln Pro Asp Trp Ser Glu Ala Phe Gln Cys
 20 25 30
 Val Ser Ala Val Glu Val Glu Ile His Ala Gly Gly Leu Asn Leu Glu
 35 40 45
 Pro Phe Leu Lys Arg Val Asp Ala Asp Ala Asp Pro Arg Gln Tyr Ala
 50 55 60
 Asp Thr Val Lys Ala Leu Arg Val Arg Arg Leu Thr Val Gly Ala Ala
 65 70 75 80
 Gln Val Pro Ala Gln Leu Leu Val Gly Ala Leu Arg Val Leu Ala Tyr
 85 90 95
 Ser Arg Leu Lys Glu Leu Thr Leu Glu Asp Leu Lys Ile Thr Gly Thr
 100 105 110
 Met Pro Pro Leu Pro Leu Glu Ala Thr Gly Leu Ala Leu Ser Ser Leu
 115 120 125
 Arg Leu Arg Asn Val Ser Trp Ala Thr Gly Arg Ser Trp Leu Ala Glu
 130 135 140
 Leu Gln Gln Trp Leu Lys Pro Gly
 145 150

(2) INFORMATION FOR SEQ ID NO:15:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 138 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

Cys Val Cys Asn Phe Ser Glu Pro Gln Pro Asp Trp Ser Glu Ala Phe
 1 5 10 15
 Gln Cys Val Ser Ala Val Glu Val Glu Ile His Ala Gly Gly Leu Asn
 20 25 30

Leu Glu Pro Phe Leu Lys Arg Val Asp Ala Asp Ala Asp Pro Arg Gln
 35 40 45
 Tyr Ala Asp Thr Val Lys Ala Leu Arg Val Arg Arg Leu Thr Val Gly
 50 55 60
 Ala Ala Gln Val Pro Ala Gln Leu Leu Val Gly Ala Leu Arg Val Leu
 65 70 75 80
 Ala Tyr Ser Arg Leu Lys Glu Leu Thr Leu Glu Asp Leu Lys Ile Thr
 85 90 95
 Gly Thr Met Pro Pro Leu Pro Leu Glu Ala Thr Gly Leu Ala Leu Ser
 100 105 110
 Ser Leu Arg Leu Arg Asn Val Ser Trp Ala Thr Gly Arg Ser Trp Leu
 115 120 125
 Ala Glu Leu Gln Gln Trp Leu Lys Pro Gly
 130 135

(2) INFORMATION FOR SEQ ID NO:16:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 41 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

CGCCAGAACC TTGTGCAGCT GCCGCTGAAG ATTTCCGCTG C

(2) INFORMATION FOR SEQ ID NO:17:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 41 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

GTGAGCTGGA CGATGCAGCT GCCGCCTGCG TCTGCAACTT C

(2) INFORMATION FOR SEQ ID NO:18:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 40 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

CCGCTGCGTC TCGCAGCTG CCGCACCTCA GCCCGACTGG

(2) INFORMATION FOR SEQ ID NO:19:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 41 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:

GCAACTTCTC CGAAGCAGCT GCCGCCTGGT CCGAAGCCTT C

(2) INFORMATION FOR SEQ ID NO:20:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 39 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:

GAACTCAGC CCGACGCAGC TGCAGCCTTC CAGTGTGTG

(2) INFORMATION FOR SEQ ID NO:21:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 41 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:21:

CCGACTGGTC CGAAGCAGCT GCGTGTGTGT CTGCAGTAGA G

(2) INFORMATION FOR SEQ ID NO:22:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 40 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:22:

CATGCCGCG GTGCAGCTGC AGCGCCGTTT CTAAAGCGCG

(2) INFORMATION FOR SEQ ID NO:23:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 42 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:23:

GGTCTCAACC TAGAGGCAGC TGCAGCGCGC GTCGATGCGG AC

(2) INFORMATION FOR SEQ ID NO:24:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 39 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:24:

GAGCCGTTTC TAAAGGCAGC TGCTGCGGAC GCCGACCCG

(2) INFORMATION FOR SEQ ID NO:25:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 33 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:25:

CATGGAGGGA CTTCCGCTG GGGACTTTC AGC

(2) INFORMATION FOR SEQ ID NO:26:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 33 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:26:

CATGGCTGGA AAGTCCCCAG CGGAAAGTCC CTC

(2) INFORMATION FOR SEQ ID NO:27:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 348 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:27:

Thr	Thr	Pro	Glu	Pro	Cys	Glu	Leu	Asp	Asp	Glu	Asp	Phe	Arg	Cys	Val
1				5					10					15	

Cys Asn Phe Ser Glu Pro Gln Pro Asp Trp Ser Glu Ala Phe Gln Cys
 20 25 30
 Val Ser Ala Val Glu Val Glu Ile His Ala Gly Gly Leu Asn Leu Glu
 35 40 45
 Pro Phe Leu Lys Arg Val Asp Ala Asp Ala Asp Pro Arg Gln Tyr Ala
 50 55 60
 Asp Thr Val Lys Ala Leu Arg Val Arg Arg Leu Thr Val Gly Ala Ala
 65 70 75 80
 Gln Val Pro Ala Gln Leu Leu Val Gly Ala Leu Arg Val Leu Ala Tyr
 85 90 95
 Ser Arg Leu Lys Glu Leu Thr Leu Glu Asp Leu Lys Ile Thr Gly Thr
 100 105 110
 Met Pro Pro Leu Pro Leu Glu Ala Thr Gly Leu Ala Leu Ser Ser Leu
 115 120 125
 Arg Leu Arg Asn Val Ser Trp Ala Thr Gly Arg Ser Trp Leu Ala Glu
 130 135 140
 Leu Gln Gln Trp Leu Lys Pro Gly Leu Lys Val Leu Ser Ile Ala Gln
 145 150 155 160
 Ala His Ser Pro Ala Phe Ser Cys Glu Gln Val Arg Ala Phe Pro Ala
 165 170 175
 Leu Thr Ser Leu Asp Leu Ser Asp Asn Pro Gly Leu Gly Glu Arg Gly
 180 185 190
 Leu Met Ala Ala Leu Cys Pro His Lys Phe Pro Ala Ile Gln Asn Leu
 195 200 205
 Ala Leu Arg Asn Thr Gly Met Glu Thr Pro Thr Gly Val Cys Ala Ala
 210 215 220
 Leu Ala Ala Ala Gly Val Gln Pro His Ser Leu Asp Leu Ser His Asn
 225 230 235 240
 Ser Leu Arg Ala Thr Val Asn Pro Ser Ala Pro Arg Cys Met Trp Ser
 245 250 255
 Ser Ala Leu Asn Ser Leu Asn Leu Ser Phe Ala Gly Leu Glu Gln Val
 260 265 270
 Pro Lys Gly Leu Pro Ala Lys Leu Arg Val Leu Asp Leu Ser Cys Asn
 275 280 285
 Arg Leu Asn Arg Ala Pro Gln Pro Asp Glu Leu Pro Glu Val Asp Asn

290		295		300
Leu Thr Leu Asp Gly Asn Pro Phe Leu Val Pro Gly Thr Ala Leu Pro				
305		310		315 320
His Glu Gly Ser Met Asn Ser Gly Val Val Pro Ala Cys Ala Arg Ser				
		325		330 335
Thr Leu Ser Val Gly Val Ser Gly Thr Leu Val Leu				
		340		345

(2) INFORMATION FOR SEQ ID NO:28:

(i) SEQUENCE CHARACTERISTICS:

- SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 60 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:28:

```

11) SEQUENCE DESCRIPTION:
Thr Thr Pro Glu Pro Cys Glu Leu Asp Asp Glu Asp Phe Arg Cys Val
1          5          10          15
Cys Asn Phe Ser Glu Pro Gln Pro Asp Trp Ser Glu Ala Phe Gln Cys
          20          25          30
Val Ser Ala Val Glu Val Glu Ile His Ala Gly Gly Leu Asn Leu Glu
          35          40          45
Pro Phe Leu Lys Arg Val Asp Ala Asp Ala Asp Pro
          50          55          60

```

(2) INFORMATION FOR SEQ ID NO:29:

(i) SEQUENCE CHARACTERISTICS:

- SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 60 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:29:

xi) SEQUENCE DESCRIPTION: 321

Thr Thr Pro Glu Pro Cys Ala Ala Ala Ala Glu Asp Phe Arg Cys Val
1 5 10 15

- 59 -

Cys Asn Phe Ser Glu Pro Gln Pro Asp Trp Ser Glu Ala Phe Gln Cys
 20 25 30
 Val Ser Ala Val Glu Val Glu Ile His Ala Gly Gly Leu Asn Leu Glu
 35 40 45
 Pro Phe Leu Lys Arg Val Asp Ala Asp Ala Asp Pro
 50 55 60

(2) INFORMATION FOR SEQ ID NO:30:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 60 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:30:

Thr Thr Pro Glu Pro Cys Glu Leu Asp Asp Ala Ala Ala Ala Cys Val
 1 5 10 15
 Cys Asn Phe Ser Glu Pro Gln Pro Asp Trp Ser Glu Ala Phe Gln Cys
 20 25 30
 Val Ser Ala Val Glu Val Glu Ile His Ala Gly Gly Leu Asn Leu Glu
 35 40 45
 Pro Phe Leu Lys Arg Val Asp Ala Asp Ala Asp Pro
 50 55 60

(2) INFORMATION FOR SEQ ID NO:31:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 60 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:31:

Thr Thr Pro Glu Pro Cys Glu Leu Asp Asp Glu Asp Phe Arg Cys Val
 1 5 10 15

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 60 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:32:

```

xi) SEQUENCE DESCRIPTION: 121
  Thr Thr Pro Glu Pro Cys Glu Leu Asp Asp Glu Asp Phe Arg Cys Val
  1      5      10      15
  Cys Asn Phe Ser Glu Ala Ala Ala Ala Trp Ser Glu Ala Phe Gln Cys
      20      25      30
  Val Ser Ala Val Glu Val Glu Ile His Ala Gly Gly Leu Asn Leu Glu
      35      40      45
  Pro Phe Leu Lys Arg Val Asp Ala Asp Ala Asp Pro
      50      55      60

```

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 60 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:33:

(xi) SEQUENCE DESCRIPTION: SEQ ID: 1

Thr	Thr	Pro	Glu	Pro	Cys	Glu	Leu	Asp	Asp	Glu	Asp	Phe	Arg	Cys	Val
1				5					10					15	
Cys	Asn	Phe	Ser	Glu	Pro	Gln	Pro	Asp	Ala	Ala	Ala	Ala	Phe	Gln	Cys
			20					25					30		

Val Ser Ala Val Glu Val Glu Ile His Ala Gly Gly Leu Asn Leu Glu
 35 40 45
 Pro Phe Leu Lys Arg Val Asp Ala Asp Ala Asp Pro
 50 55 60

(2) INFORMATION FOR SEQ ID NO:34:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 60 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:34:

Thr Thr Pro Glu Pro Cys Glu Leu Asp Asp Glu Asp Phe Arg Cys Val
 1 5 10 15
 Cys Asn Phe Ser Glu Pro Gln Pro Asp Trp Ser Glu Ala Ala Ala Cys
 20 25 30
 Val Ser Ala Val Glu Val Glu Ile His Ala Gly Gly Leu Asn Leu Glu
 35 40 45
 Pro Phe Leu Lys Arg Val Asp Ala Asp Ala Asp Pro
 50 55 60

(2) INFORMATION FOR SEQ ID NO:35:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 60 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:35:

Thr Thr Pro Glu Pro Cys Glu Leu Asp Asp Glu Asp Phe Arg Cys Val
 1 5 10 15
 Cys Asn Phe Ser Glu Pro Gln Pro Asp Trp Ser Glu Ala Phe Gln Cys
 20 25 30
 Val Ser Ala Val Glu Val Glu Ile His Ala Gly Gly Ala Ala Ala Ala

35 40 45
 Pro Phe Leu Lys Arg Val Asp Ala Asp Ala Asp Pro
 50 55 60

(2) INFORMATION FOR SEQ ID NO:36:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 60 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:36:

Thr Thr Pro Glu Pro Cys Glu Leu Asp Asp Glu Asp Phe Arg Cys Val
 1 5 10 15
 Cys Asn Phe Ser Glu Pro Gln Pro Asp Trp Ser Glu Ala Phe Gln Cys
 20 25 30
 Val Ser Ala Val Glu Val Glu Ile His Ala Gly Gly Leu Asn Leu Glu
 35 40 45
 Ala Ala Ala Ala Arg Val Asp Ala Asp Ala Asp Pro
 50 55 60

(2) INFORMATION FOR SEQ ID NO:37:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 60 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:37:

Thr Thr Pro Glu Pro Cys Glu Leu Asp Asp Glu Asp Phe Arg Cys Val
 1 5 10 15
 Cys Asn Phe Ser Glu Pro Gln Pro Asp Trp Ser Glu Ala Phe Gln Cys
 20 25 30
 Val Ser Ala Val Glu Val Glu Ile His Ala Gly Gly Leu Asn Leu Glu
 35 40 45
 Pro Phe Leu Lys Ala Ala Ala Ala Asp Ala Asp Pro
 50 55 60

- 63 -

The embodiments of the invention in which an exclusive property or privilege is claimed are defined as follows:

5

1. A polypeptide comprising an amino acid sequence that begins with one of amino acids 1 through 6 and ends with one of amino acids 152 through 348 of the sequence of FIG. 1, wherein X_1 - X_4 are each independently selected from Gly and amino acids that have a C_1 - C_6 aliphatic side chain; and physiologically acceptable salts thereof.

2. A polypeptide according to Claim 1, wherein said amino acid sequence comprises amino acids 1-152 of the sequence of FIG. 1.

3. A polypeptide according to Claim 1, wherein said amino acid sequence comprises amino acids 1-348 of the sequence of FIG. 1.

4. A polypeptide according to Claim 1, wherein X_1 - X_4 are each independently selected from Gly, Ala, Val, Leu, Ile and Pro.

25

5. A polypeptide according to Claim 1, wherein X_1 - X_4 are each Ala.

6. A polypeptide according to Claim 1, wherein X_1 - X_4 are each amino acids of the L configuration.

7. A polypeptide according to Claim 1, wherein said sequence comprises amino acids 1-152 or 1-348 of the sequence of FIG. 1 and wherein X_1 - X_4 are each L-Ala.

- 64 -

8. A polypeptide comprising an amino acid sequence that begins with amino acid 15 and ends with one of amino acids 152 through 348 of the sequence of FIG.1, and physiologically acceptable salts thereof.

9. A polypeptide according to Claim 8, wherein said amino acid sequence comprises amino acids 15-152 of the sequence of FIG. 1.

10

10. A polypeptide according to Claim 8, wherein said amino acid sequence comprises amino acids 15-348 of the sequence of FIG. 1.

15

11. A polynucleotide that encodes a polypeptide according to Claim 8.

20

12. A polynucleotide according to Claim 11 which is DNA.

25


13. A method of treating an inflammatory condition in a patient in need thereof, which comprises administering to said patient an effective amount of a polypeptide according to Claim 1 or Claim 8.

30

14. A method according to Claim 13 wherein said amount is from 0.1 mg/kg to 100 mg/kg.

15. A pharmaceutical composition comprising a peptide according to Claim 1 or Claim 8 in admixture with a pharmaceutically acceptable carrier thereof.

Figure 1 : CD14 Disulfide Structure 50



Sequence and expression of SCD14 alanine substitution mutants.

Name	Sequence of mutant SCD14	Expression in COS-7 cells
1. SCD14 ₁₋₃₄₆	TTPEPCELDEDFRCVCFSEFPQPDWSEAFQCVSAREVEHAGGLNEPFLKRVADADP	+
2. SCD14 _{17-109A}	AAAA	+
3. SCD14 _{111-149A}	AAAA	+
4. SCD14 _{110-219A}	AAAA	+
5. SCD14 _{122-259A}	AAAA	+
6. SCD14 _{120-299A}	AAA	+
7. SCD14 _{120-319A}	AA	+
8. SCD14 _{105-409A}	AAAA	+
9. SCD14 _{100-529A}	AAAA	+
10. SCD14 _{153-559A}	AAA	+

1 Sequence between amino acids 1 and 60 from mature SCD14 are shown. Solid lines indicate the same sequences as SCD14₁₋₃₄₆.

2 Expression of SCD14 proteins in COS-7 cells was determined by Western blot using polyclonal anti-CD14 antiserum

FIGURE 2




FIGURE 3




FIGURE 4





FIGURE 5A






FIGURE 6




FIGURE 7A




FIGURE 7B




FIGURE 8A




FIGURE 8B




FIGURE 9

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 95/17095

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 C07K14/705 C12N15/12 A61K38/17

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 6 C07K C12N A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO,A,93 19772 (NORTH SHORE UNIVERSITY HOSPITA) 14 October 1993 see the whole document ---	8-15
P,X	JOURNAL OF BIOLOGICAL CHEMISTRY, vol. 270, no. 29, 21 July 1995, MD US, pages 17237-17242, XP002003900 T.S.-C.JUAN E.A.: "Identification of a domain in sCD14 essential for LPS signaling but not LPS binding" see the whole document ---	1-7, 13-15
A	WO,A,91 01639 (SCRIPPS CLINIC RES ;UNIV ROCKEFELLER (US)) 21 February 1991 see page 16, line 32 - page 17, line 12; claims 18,19 -----	1-15

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- * 'A' document defining the general state of the art which is not considered to be of particular relevance
- * 'E' earlier document but published on or after the international filing date
- * 'L' document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- * 'O' document referring to an oral disclosure, use, exhibition or other means
- * 'P' document published prior to the international filing date but later than the priority date claimed

* 'T' later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

* 'X' document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

* 'Y' document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

* 'a' document member of the same patent family

Date of the actual completion of the international search

24 May 1996

Date of mailing of the international search report

12.06.96

Name and mailing address of the ISA
European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+31-70) 340-3016

Authorized officer

Groenendijk, M

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/JS 95/17095

Patent document cited in search report	Publication date	Patent family member(s)		Publication date
WO-A-9319772	14-10-93	CA-A-	2133758	14-10-93
		EP-A-	0634935	25-01-95
WO-A-9101639	21-02-91	AU-B-	645515	20-01-94
		AU-B-	6146190	11-03-91
		CA-A-	2022429	02-02-91
		EP-A-	0485430	20-05-92
		GR-A-	90100582	30-12-91
		JP-T-	5501399	18-03-93

Form PCT/ISA/210 (patent family annex) (July 1992)